

p-(SULFONYL) ARYL AND HETEROARYLAMINES-
AS ANTI-INFLAMMATORY AGENTS

Patent Application
Attorney Docket No. R0070B-REG

Nancy Elisabeth Krauss
Tara Mirzadegan
David Bernard Smith
Keith Adrian Walker

CERTIFICATE OF EXPRESS MAILING

I hereby certify that this correspondence and patent application are being deposited with the U.S. Postal Service as "EXPRESS MAIL - POST OFFICE TO ADDRESSEE" under 37 CFR 1.10 in an envelope addressed to: Assistant Commissioner of Patents, Washington D.C. 20231 on APRIL 26, 2001.

EXPRESS MAIL Mailing Label No **EH814670585US**

Name of Person Mailing Chris Myers

Signature Chris Myers Date April 26, 2001

BACKGROUND OF THE INVENTION

Cross Reference to Related Applications

Pursuant to 35 U.S.C. §119(e), this application claims priority to the filing date of the
5 United States Provisional Patent Application Serial No. 60/200,310, the disclosure of which is
herein incorporated by reference.

Field of the Invention

This invention relates to anti-inflammatory and analgesic compounds, especially to
10 certain *p*-(sulfonyl)-aryl and -heteroaryl amines, pharmaceutical compositions containing them,
methods for their use, and methods for preparing these compounds.

Description of the Related Art

U.S. Patent No. 4,277,492 (Dow Chemical) discloses 4-
15 bis((phenylmethyl)amino)benzenesulfonic acids and their pharmaceutically-acceptable salts useful
as antiviral agents.

U.S. Patent No. 4,857,530 (Warner-Lambert) discloses 6-substituted-4(3H)-
quinazolinones which inhibit the enzyme thymidylate synthase and thus are useful as anticancer
agents.

20 U.S. Patent No. 5,538,976 (Yamanouchi Pharmaceutical) discloses substituted tertiary
amino compounds, one of the substituents being a pyrimidine ring, a pyridazine ring, or a triazine
ring, or pharmaceutically acceptable salts thereof. These compounds have aromatase inhibiting
activity and are useful as a prophylactics and/or therapeutic agents for breast cancer, mastopathy,
endometriosis, prostatic-hypertrophy, and so forth.


25 WO 98/25893 (Athena Neurosciences) discloses arylsulfonamides which have activity as
inhibitors of phospholipase A₂, inhibitors of cytokine release, and as inhibitors of
neurodegeneration.

WO 98/50029 (University of Pittsburgh) discloses certain substituted
benzenesulfonamides as inhibitors of protein isoprenyl transferases.

EP 757037 A2 (Ono Pharmaceutical) discloses certain benzenesulfonyl amino acids as metalloproteinase inhibitors.

SUMMARY OF THE INVENTION

5 In a first aspect, this invention provides compounds selected from the group of compounds represented by formula (I):



10 Formula I

wherein:

A is $-(CR_2)_n-$ where n is 1, 2, or 3 and R is independently hydrogen or alkyl;

B is aryl or heteroaryl;

X and Y are, independently, CH or nitrogen;

15 R¹ is alkyl, alkenyl, cyanoalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heteroaralkyl, heterocyclyl, heterocyclylalkyl, heteroalkyl or alkylcarbonylalkyl;

R² is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkenyl, hydroxyalkyl, alkoxyalkyl, alkoxyalkyl, aryl, aralkyl, or NR¹³R¹⁴ wherein

R¹³ is hydrogen or alkyl;

20 R¹⁴ is hydrogen, alkyl, alkenyl, acyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aralkyl, hydroxyalkyl, alkoxyalkyl, carboxyalkyl, alkoxyalkyl, or aminoalkyl;

R³ is hydrogen, alkyl, halo, nitro, cyano, hydroxy or alkoxy; and

prodrugs, individual isomers, mixtures of isomers, and pharmaceutically acceptable salts thereof.

In a second aspect, this invention provides pharmaceutical compositions containing a therapeutically effective amount of a compound of formula (I) or its pharmaceutically acceptable salt and a pharmaceutically acceptable excipient.

In a third aspect, this invention provides a method of treatment of a disease, in particular an inflammatory or autoimmune disease, in a mammal treatable by administration of a prostaglandin G/H synthase inhibitor, comprising administration of a therapeutically effective amount of a compound of formula (I) or its pharmaceutically acceptable salt.

In a fourth aspect, this invention provides processes for preparing compounds of formula (I).

DETAILED DESCRIPTION OF THE INVENTION

Definitions

Unless otherwise stated, the following terms used in the specification and claims have the meanings given below:

"Acyl" means the group $-C(O)R'$, where R' is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, phenyl or phenylalkyl.

"Alkyl" means a linear saturated monovalent hydrocarbon radical of one to six carbon atoms or a branched saturated monovalent hydrocarbon radical of three to six carbon atoms, *e.g.*, methyl, ethyl, *n*-propyl, 2-propyl, *tert*-butyl, pentyl, and the like.

"Alkylene" means a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms, *e.g.*, methylene, ethylene, propylene, 2-methylpropylene, pentylene, and the like.

"Alkenyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, *e.g.*, ethenyl, propenyl, and the like.

"Alkynyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms, containing at least one triple bond, *e.g.*, ethynyl, propynyl, and the like.

"Alkoxy", "aryloxy", "aralkyloxy", or "heteroaralkyloxy" means a radical -OR where R is an alkyl, aryl, aralkyl, or heteroaralkyl respectively, as defined herein, *e.g.*, methoxy, phenoxy, benzyloxy, pyridin-2-ylmethoxy, and the like.

"Alkoxycarbonylalkyl" means a radical $-R^aC(O)R^b$ where R^a is an alkylene group as defined above and R^b is an alkoxy group as defined above *e.g.*, methoxycarbonylethyl, ethoxycarbonylbutyl, and the like.

"Aryl" means a monovalent monocyclic or bicyclic aromatic hydrocarbon radical of 6 to 10 ring atoms which is substituted independently with one to four substituents, preferably one, two, or three substituents selected from alkyl, cycloalkyl, cycloalkyl-alkyl, halo, nitro, cyano, hydroxy, alkoxy, alkylthio, alkylsulfonyl, amino, acylamino, mono-alkylamino, di-alkylamino, haloalkyl, haloalkoxy, heteroalkyl, COR (where R is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, phenyl or phenylalkyl), $-(CR'R'')_n-COOR$ (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R is hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl) or $-(CR'R'')_n-CONR^aR^b$ (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R^a and R^b are, independently of each other, hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl). More specifically the term aryl includes, but is not limited to, phenyl, biphenyl, 1-naphthyl, and 2-naphthyl, and the derivatives thereof.

"Aralkyl" means a radical $-R^aR^b$ where R^a is an alkylene group and R^b is an aryl group as defined herein, *e.g.*, benzyl, phenylethyl, 3-(3-chlorophenyl)-2-methylpentyl, and the like.

"Aralkenyl" means a radical $-R^aR^b$ where R^a is an alkenylene group and R^b is an aryl group as defined herein, *e.g.*, 3-phenyl-2-propenyl, and the like.

"Arylheteroalkyl" means a radical $-R^aR^b$ where R^a is an heteroalkylene group and R^b is an aryl group as defined herein, *e.g.*, 2-hydroxy-2-phenyl-ethyl, 2-hydroxy-1-hydroxymethyl-2-phenyl-ethyl, and the like.

"Cycloalkyl" means a saturated monovalent cyclic hydrocarbon radical of three to seven ring carbons. The cycloalkyl may be optionally substituted independently with one, two, or three substituents selected from alkyl, optionally substituted phenyl, or $-C(O)R$ (where R is hydrogen, alkyl, haloalkyl, amino, acylamino, mono-alkylamino, di-alkylamino, hydroxy, alkoxy, or

optionally substituted phenyl). More specifically, the term cycloalkyl includes, for example, cyclopropyl, cyclohexyl, phenylcyclohexyl, 4-carboxycyclohexyl, 2-carboxamidocyclohexyl, 2-dimethylaminocarbonyl-cyclohexyl, and the like.

"Cycloalkyl-alkyl" means a radical $-R^aR^b$ where R^a is an alkylene group and R^b is a cycloalkyl group as defined herein, *e.g.*, cyclopropylmethyl, cyclohexylpropyl, 3-cyclohexyl-2-methylpropyl, and the like.

"Haloalkyl" means alkyl substituted with one or more same or different halo atoms, *e.g.*, $-\text{CH}_2\text{Cl}$, $-\text{CF}_3$, $-\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CCl}_3$, and the like, and further includes those alkyl groups such as perfluoroalkyl in which all hydrogen atoms are replaced by fluorine atoms.

"Heteroalkyl" means an alkyl radical as defined herein with one, two or three substituents independently selected from $-\text{OR}^a$, $-\text{NR}^b\text{R}^c$, and $-\text{S}(\text{O})_n\text{R}^d$ (where n is an integer from 0 to 2), with the understanding that the point of attachment of the heteroalkyl radical is through a carbon atom of the heteroalkyl radical. R^a is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, aryl, aralkyl, alkoxycarbonyl, aryloxy carbonyl, carboxamido, or mono- or di-alkylcarbamoyl. R^b is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, aryl or aralkyl. R^c is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, aryl, aralkyl, acyl, alkoxycarbonyl, aryloxy carbonyl, carboxamido, mono- or di-alkylcarbamoyl or alkylsulfonyl. R^d is hydrogen (provided that n is 0), alkyl, cycloalkyl, cycloalkyl-alkyl, aryl, aralkyl, heteroaryl, amino, mono-alkylamino, di-alkylamino, hydroxyalkyl or hydroxyalkylamino. Representative examples include, for example, 2-hydroxyethyl, 2,3-dihydroxypropyl, 2-methoxyethyl, benzyloxymethyl, 2-methylsulfonyl-ethyl.

"Heteroaryl" means a monovalent monocyclic or bicyclic radical of 5 to 12 ring atoms having at least one aromatic ring containing one, two, or three ring heteroatoms selected from N, O, or S, the remaining ring atoms being C, with the understanding that the attachment point of the heteroaryl radical will be on an aromatic ring. The heteroaryl ring is optionally substituted independently with one to four substituents, preferably one or two substituents, selected from alkyl, cycloalkyl, cycloalkyl-alkyl, halo, nitro, cyano, hydroxy, alkoxy, amino, acylamino, mono-alkylamino, di-alkylamino, haloalkyl, haloalkoxy, heteroalkyl, $-\text{COR}$ (where R is hydrogen, alkyl, phenyl or phenylalkyl, $-(\text{CR}'\text{R}'')_n-\text{COOR}$ (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, phenyl or

phenylalkyl), or $-(CR'R'')_n-CONR^aR^b$ (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R^a and R^b are, independently of each other, hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, phenyl or phenylalkyl). More specifically the term heteroaryl includes, but is not limited to, pyridyl, furanyl, thienyl, thiazolyl, isothiazolyl, triazolyl, imidazolyl, isoxazolyl, pyrrolyl, pyrazolyl, pyridazinyl, pyrimidinyl, benzofuranyl, tetrahydrobenzofuranyl, isobenzofuranyl, benzothiazolyl, benzoisothiazolyl, benzotriazolyl, indolyl, isoindolyl, benzoxazolyl, quinolyl, tetrahydroquinolyl, isoquinolyl, benzimidazolyl, benzisoxazolyl or benzothienyl, and the derivatives thereof.

"Heteroaralkyl" means a radical $-R^aR^b$ where R^a is an alkylene group and R^b is a heteroaryl group as defined herein, *e.g.*, pyridin-3-ylmethyl, 3-(benzofuran-2-yl)propyl, and the like.

"Heteroaralkenyl" means a radical $-R^aR^b$ where R^a is an alkenylene group and R^b is a heteroaryl group as defined herein, *e.g.*, 3-(pyridin-3-yl)propen-2-yl, and the like.

"Heterocyclyl" means a saturated or unsaturated non-aromatic cyclic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatoms selected from O, NR (where R is independently hydrogen or alkyl or S(O)_n (where n is an integer from 0 to 2), the remaining ring atoms being C, where one or two C atoms may optionally be replaced by a carbonyl group. The heterocyclyl ring may be optionally substituted independently with one, two, or three substituents selected from alkyl, cycloalkyl, cycloalkyl-alkyl, halo, nitro, cyano, hydroxy, alkoxy, amino, mono-alkylamino, di-alkylamino, haloalkyl, haloalkoxy, -COR (where R is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, phenyl or phenylalkyl), $-(CR'R'')_n-COOR$ (n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, phenyl or phenylalkyl), or $-(CR'R'')_n-CONR^aR^b$ (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R^a and R^b are, independently of each other, hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl). More specifically the term heterocyclyl includes, but is not limited to, tetrahydropyranyl, piperidino, N-methylpiperidin-3-yl, piperazino, N-methylpyrrolidin-3-yl, 3-pyrrolidino, 2-pyrrolidon-1-yl, morpholino, thiomorpholino, thiomorpholino-1-oxide, thiomorpholino-1,1-dioxide, pyrrolidinyl, and the derivatives thereof.

"Heterocyclalkyl" means a radical $-R^aR^b$ where R^a is an alkylene group and R^b is a heterocycl group as defined herein, *e.g.*, tetrahydropyran-2-ylmethyl, 4-methylpiperazin-1-ylethyl, 3-piperidinylmethyl, and the like.

"Heteroalkylene" means a linear saturated divalent hydrocarbon radical of one to six carbons or a branched saturated hydrocarbon radical of three to six carbon atoms with one, two or three substituents independently selected from $-OR^a$, $-NR^bR^c$, and $-S(O)_nR^d$ (where n is an integer from 0 to 2) where, R^a , R^b , R^c , and R^d are as defined herein for a heteroalkyl radical. Examples include, 2-hydroxyethan-1,2-diyl, 2-hydroxypropan-1,3-diyl and the like.

"Heterosubstituted cycloalkyl" means a cycloalkyl group wherein one, two, or three hydrogen atoms are replaced by substituents independently selected from the group consisting of hydroxy, alkoxy, amino, acylamino, mono-alkylamino, di-alkylamino, or $-SO_nR$ (where n is an integer from 0 to 2 and when n is 0, R is hydrogen or alkyl and when n is 1 or 2, R is alkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heteroaryl, amino, acylamino, mon-alkylamino, di-alkylamino, or hydroxyalkyl). Examples include 4-hydroxycyclohexyl, 2-aminocyclohexyl etc.

"Heteroalkylsubstituted cycloalkyl" means a cycloalkyl group wherein one, two, or three hydrogen atoms are replaced independently by heteroalkyl groups, with the understanding that the heteroalkyl group is attached to the cycloalkyl group via a carbon-carbon bond. Examples include 1-hydroxymethyl-cyclopent-1-yl, 2-hydroxymethyl-cyclohex-2-yl and the like.

"Heteroalkylsubstituted heterocycl" means a heterocycl group wherein one, two, or three hydrogen atoms are replaced independently by heteroalkyl groups, with the understanding that the heteroalkyl group is attached to the heterocycl group via a carbon-carbon bond. Examples include 4-hydroxymethyl-piperidin-1-yl, and the like.

"Hydroxyalkyl" means an alkyl radical as defined herein, substituted with one or more, preferably one, two or three hydroxy groups, provided that the same carbon atom does not carry more than one hydroxy group. Representative examples include, but are not limited to, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 1-hydroxymethyl-2-methylpropyl, 2-hydroxybutyl, 3-hydroxybutyl, 4-hydroxybutyl, 2,3-dihydroxypropyl, 1-hydroxymethyl-2-hydroxyethyl, 2,3-dihydroxybutyl, 3,4-dihydroxybutyl and 2-hydroxymethyl-3-hydroxypropyl, preferably 2-hydroxyethyl, 2,3-dihydroxypropyl and 1-hydroxymethyl-2-hydroxyethyl.

Accordingly, as used herein, the term "hydroxyalkyl" is used to define a subset of heteroalkyl groups.

"Optionally substituted phenyl" means a phenyl ring which is optionally substituted independently with one to four substituents, preferably one or two substituents selected from alkyl, cycloalkyl, cycloalkyl-alkyl, halo, nitro, cyano, hydroxy, alkoxy, amino, acylamino, mono-alkylamino, di-alkylamino, haloalkyl, haloalkoxy, heteroalkyl, -COR (where R is hydrogen, alkyl, phenyl or phenylalkyl, -(CR'R'')_n-COOR (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R is hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl), or -(CR'R'')_n-CONR^aR^b (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R^a and R^b are, independently of each other, hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl).

"Leaving group" has the meaning conventionally associated with it in synthetic organic chemistry i.e., an atom or group capable of being displaced by a nucleophile and includes halo (such as chloro, bromo, iodo), alkanesulfonyloxy, arenesulfonyloxy, alkylcarbonyloxy (e.g. acetoxy), arylcarbonyloxy, mesyloxy, tosyloxy, trifluoromethanesulfonyloxy, aryloxy (e.g., 2,4-dinitrophenoxy), methoxy, N,O-dimethylhydroxylamino, and the like.

"Pharmaceutically acceptable excipient" means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes an excipient that is acceptable for veterinary use as well as human pharmaceutical use. A "pharmaceutically acceptable excipient" as used in the specification and claims includes both one and more than one such excipient.

"Pharmaceutically acceptable salt" of a compound means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. Such salts include:

(1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid,

2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid, glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or

(2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, tromethamine, *N*-methylglucamine, and the like.

"Pro-drugs" means any compound which releases an active parent drug according to Formula (I) *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of Formula (I) are prepared by modifying functional groups present in the compound of Formula (I) in such a way that the modifications may be cleaved *in vivo* to release the parent compound. Prodrugs include compounds of Formula (I) wherein a hydroxy, amino, or sulfhydryl group in a compound of Formula (I) is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino, or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to esters (e.g., acetate, formate, and benzoate derivatives), carbamates (e.g., *N,N*-dimethylaminocarbonyl) of hydroxy functional groups in compounds of Formula (I), and the like.

"Protecting group" refers to a grouping of atoms that when attached to a reactive group in a molecule masks, reduces or prevents that reactivity. Examples of protecting groups can be found in T.W. Greene and P.G. Futs, Protective Groups in Organic Chemistry, (Wiley, 2nd ed. 1991) and Harrison and Harrison *et al.*, Compendium of Synthetic Organic Methods, Vols. 1-8 (John Wiley and Sons. 1971-1996). Representative amino protecting groups include formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (CBZ), *tert*-butoxycarbonyl (Boc), trimethylsilyl (TMS), 2-trimethylsilyl-ethanesulfonyl (SES), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (Fmoc), nitro-veratryloxycarbonyl (NVOC) and the like. Representative hydroxy protecting groups include those where the hydroxy group is either acylated or alkylated such as benzyl and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers and allyl ethers.

"Treating" or "treatment" of a disease includes:

(1) preventing the disease, i.e. causing the clinical symptoms of the disease not to develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease,

5 (2) inhibiting the disease, i.e., arresting or reducing the development of the disease or its clinical symptoms, or

(3) relieving the disease, i.e., causing regression of the disease or its clinical symptoms.

"A therapeutically effective amount" means the amount of a compound that, when
10 administered to a mammal for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

"Optional" or "optionally" in the above definitions means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where
15 the event or circumstance occurs and instances in which it does not. For example, "heterocyclo group optionally mono- or di- substituted with an alkyl group" means that the alkyl may but need not be present, and the description includes situations where the heterocyclo group is mono- or disubstituted with an alkyl group and situations where the heterocyclo group is not substituted with the alkyl group.


20 Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers". Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers" and those that are non-superimposable mirror images of each other are termed "enantiomers". When a compound has an
25 asymmetric center, for example, it is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the *R*- and *S*-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral compound can exist as either

individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a “racemic mixture”.

The compounds of this invention may exist in stereoisomeric form if they possess one or more asymmetric centers or a double bond with asymmetric substitution and, therefore, can be produced as individual stereoisomers or as mixtures. Unless otherwise indicated, the description is intended to include individual stereoisomers as well as mixtures. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art (see discussion in Chapter 4 of “Advanced Organic Chemistry”, 4th edition J. March, John Wiley and Sons, New York, 1992).

Nomenclature

The naming and numbering of the compounds of this invention is illustrated below.



(I)

The nomenclature used in this application is generally based on the IUPAC recommendations, e.g., a compound of formula (I):

where R³ is hydrogen, R² is methyl, R¹ is 2-(methylsulfonyl)ethyl, A is CH₂, B is 4-fluorophenyl, X is CH and Y is CH is named 4-{N,N-[2-(methylsulfonyl)ethyl](4-fluorobenzyl)amino}phenyl methyl sulfone.

where R³ is hydrogen, R² is methyl, R¹ is benzyl, A is CH₂, B is 4-fluorophenyl, X is CH and Y is CH is named 4-[N,N-(benzyl)(4-fluorobenzyl)amino]phenyl methyl sulfone.

where R^3 is hydrogen, R^2 is methyl, R^1 is 2-(methylsulfonyl)ethyl, A is CH_2 , B is 4-fluorophenyl, X is nitrogen and Y is CH is named 2-{N,N-[2-(methylsulfonyl)ethyl](4-fluorobenzyl)amino}pyridin-5-yl methyl sulfone.

5 where R^3 is hydrogen, R^2 is NH_2 , R^1 is 2-methylsulfonyl-ethyl, A is CH_2 , B is 4-methyl-phenyl, X is CH and Y is CH is named 4-[(2-methylsulfonyl-ethyl)-(4-methyl-benzyl)-amino]-benzenesulfonamide.


where R^3 is 3-fluoro, R^2 is methyl, R^1 is 2-methylsulfonyl-ethyl, A is CH_2 , B is 4-ethoxy-phenyl, X is CH and Y is CH is named (4-ethoxy-benzyl)-(3-fluoro-4-methanesulfonyl-phenyl)-(2-methanesulfonyl-ethyl)-amine.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172

Representative compounds of this invention are as follows:

I. Compounds of formula (I) where R² is CH₃ and the other groups are defined as below:




| CPD # | A | B | X | Y | R ¹ | R ³ | Made by Example | M. Pt. °C | M/S m/e |
|-------|-----------------|-------------------------|----|----|-------------------------|----------------|-----------------|-----------|---------|
| 1-1 | CH ₂ | phenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 367 |
| 1-2 | CH ₂ | 2-fluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 385 |
| 1-3 | CH ₂ | 2-methoxyphenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 397 |
| 1-4 | CH ₂ | 3,4-difluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 403 |
| 1-5 | CH ₂ | 2-chloro-4-fluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 419 |
| 1-6 | CH ₂ | 3,4-dimethylphenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 395 |
| 1-7 | CH ₂ | 2,4-dichlorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 436 |
| 1-8 | CH ₂ | 4-trifluoromethylphenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 435 |
| 1-9 | CH ₂ | 2-bromophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 446 |
| 1-10 | CH ₂ | 3-fluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 385 |
| 1-11 | CH ₂ | 3-chlorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 401 |

| CPD # | A | B | X | Y | R ¹ | R ³ | Made by Example | M. Pt. °C | M/S m/e |
|-------|---------------------|--------------------------------|----|----|--------------------------|----------------|-----------------|-----------|---------|
| 1-12 | CH ₂ | 4-bromophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 446 |
| 1-13 | CH ₂ | 2,3-difluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 403 |
| 1-14 | CH ₂ | 3,5-difluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 403 |
| 1-15 | CH ₂ | 4-methoxycarbonylphenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 425 |
| 1-16 | CH ₂ | 2,5-difluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 403 |
| 1-17 | CH ₂ | 2-methylthiazol-4-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 388 |
| 1-18 | CH ₂ | 4-methylthiazol-2-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 388 |
| 1-19 | CH ₂ | 2-phenylthiazol-4-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 450 |
| 1-20 | CH ₂ | 2-(4-chlorophenyl)thiazol-4-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 485 |
| 1-21 | CH ₂ | 4-fluorophenyl | CH | CH | ethyl | H | 2 | | 307 |
| 1-22 | CH ₂ | thiophen-2-yl | CH | CH | butyl | H | 3 | | 323 |
| 1-23 | CH ₂ | phenyl | CH | CH | butyl | H | 2 | | 317 |
| 1-24 | CH ₂ | 4-fluorophenyl | CH | CH | pentyl | H | 2 | | 349 |
| 1-25 | CH ₂ | 4-fluorophenyl | CH | CH | isopropyl | H | 2 | | 321 |
| 1-26 | CH ₂ | 4-fluorophenyl | CH | CH | propyl | H | E 2 | | 321 |
| 1-27 | CH ₂ | 4-fluorophenyl | CH | CH | butyl | H | 2 | | 335 |
| 1-28 | CH ₂ | 4-fluorophenyl | CH | CH | isoamyl | H | 2 | | 349 |
| 1-29 | CH ₂ | 4-fluorophenyl | CH | CH | isobutyl | H | 2 | | 335 |
| 1-30 | CH ₂ | 4-fluorophenyl | CH | CH | 2-(methoxy)ethyl | H | 2 | | 337 |
| 1-31 | (CH)CH ₃ | phenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 4 | | 381 |
| 1-32 | CH ₂ | 4-fluorophenyl | CH | CH | hydroxypropyl | H | 2 | | 337 |
| 1-33 | CH ₂ | 4-fluorophenyl | CH | CH | 3-(methylsulfonyl)propyl | H | 4 | | 399 |

| CPD # | A | B | X | Y | R ¹ | R ³ | Made by Example | M. Pt. °C | M/S m/e |
|-------|-----------------|--------------------|----|----|--|----------------|-----------------|-------------|---------|
| 1-34 | CH ₂ | 4-fluorophenyl | CH | CH | 2-(ethylsulfonyl)ethyl | H | 4 | 109.6-110.7 | 399 |
| 1-35 | CH ₂ | 4-fluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 4 | | 472 |
| 1-36 | CH ₂ | 4-fluorophenyl | CH | CH | hydroxyethyl | H | 2 | | 323 |
| 1-37 | CH ₂ | 4-fluorophenyl | CH | CH | cyclopropylmethyl | H | 2 | | 333 |
| 1-38 | CH ₂ | 4-fluorophenyl | CH | CH | 1-(tertbutyloxycarbonyl)-pyrrolidin-3-yl | H | 2 | 174.4-178.0 | 448 |
| 1-39 | CH ₂ | 4-fluorophenyl | CH | CH | 2-(morpholin-4-yl)ethyl | H | 2 | | 392 |
| 1-40 | CH ₂ | 4-fluorophenyl | CH | CH | pyrrolidin-3-yl | H | 2 | 124.0-124.3 | 348 |
| 1-41 | CH ₂ | 4-fluorophenyl | CH | CH | 3-(pyrrolidin-2-on-1-yl)-propyl | H | 2 | | 404 |
| 1-42 | CH ₂ | 4-fluorophenyl | CH | CH | benzyl | H | 1 | | 369 |
| 1-43 | CH ₂ | 4-fluorophenyl | CH | CH | 2-[N,N-(acetyl)(4-fluorobenzyl)amino]ethyl | H | 2 | | 472 |
| 1-44 | CH ₂ | 2,4-difluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 4 | 149.4-150.4 | 403 |
| 1-45 | CH ₂ | pyridin-2-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 5 | 139.5-140.5 | 368 |
| 1-46 | CH ₂ | 4-methoxyphenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 4 | | 397 |
| 1-47 | CH ₂ | 2-chlorophenyl | CH | CH | butyl | H | 2 | | 351 |
| 1-48 | CH ₂ | 2-fluorophenyl | CH | CH | butyl | H | 2 | | 335 |
| 1-49 | CH ₂ | 2-methoxyphenyl | CH | CH | butyl | H | 2 | | 347 |
| 1-50 | CH ₂ | 4-chlorophenyl | CH | CH | butyl | H | 2 | | 351 |
| 1-51 | CH ₂ | 4-methoxyphenyl | CH | CH | butyl | H | 2 | | 347 |

| CPD # | A | B | X | Y | R ¹ | R ³ | Made by Example | M. Pt. °C | M/S m/e |
|-------|---------------------------------|------------------------|----|----|-------------------------|----------------|-----------------|-------------|---------|
| 1-52 | CH ₂ | 4-cyanophenyl | CH | CH | butyl | H | 2 | | 342 |
| 1-53 | CH ₂ | 2,4-difluorophenyl | CH | CH | butyl | H | 2 | | 353 |
| 1-54 | CH ₂ | 3,4-difluorophenyl | CH | CH | butyl | H | 2 | | 353 |
| 1-55 | CH ₂ | 4-chlorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 4 | | 401 |
| 1-56 | CH ₂ | 2-chlorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 4 | | 401 |
| 1-57 | CH ₂ | 6-chloropyridazin-3-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 5 | | 403 |
| 1-58 | CH ₂ | 6-oxo-pyridazin-3-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 5 | | 385 |
| 1-59 | CH ₂ | benzothiazol-2-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 5 | | 424 |
| 1-60 | CH ₂ | 4-fluorophenyl | N | CH | 2-(methylsulfonyl)ethyl | H | 11 | | 386 |
| 1-61 | CH ₂ | 4-fluorophenyl | CH | N | 2-(methylsulfonyl)ethyl | H | 12 | | 386 |
| 1-62 | CH ₂ | thiazol-2-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 6 | | 388 |
| 1-63 | CH ₂ | 2-methoxypyridin-5-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 5 | | 398 |
| 1-64 | CH ₂ | 2-ethoxypyridin-5-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 5 | | 412 |
| 1-65 | CH ₂ | 4-ethoxyphenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 7 | | 411 |
| 1-66 | CH ₂ | pyridin-3-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 5 | | 368 |
| 1-67 | CH ₂ | pyridin-4-yl | CH | CH | 2-(methylsulfonyl)ethyl | H | 5 | 154.4-154.6 | 368 |
| 1-68 | CH ₂ | 4-fluorophenyl | CH | CH | 4-methylsulfonyl-phenyl | H | 8 | | 433 |
| 1-69 | CH ₂ | 4-fluorophenyl | CH | CH | 4-methylthio-phenyl | H | 8 | | 401 |
| 1-70 | CH ₂ | 4-fluorophenyl | CH | CH | 3-oxo-butyl | H | 9 | | 349 |
| 1-71 | (CH ₂) ₃ | phenyl | CH | CH | 2-(methylsulfonyl)ethyl | H | 4 | | 395 |
| 1-72 | CH ₂ | 4-ethoxyphenyl | CH | CH | 2-(methylsulfonyl)ethyl | 3-F | 13 | 56-59 | 429 |
| 1-73 | CH ₂ | 4-fluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | 3-F | 13 | | 403 |

II. Compounds of formula (I) where X and Y are CH, R³ is hydrogen and the other groups are defined as below:



| CPD # | A | B | X | Y | R ¹ | R ² | Made By Example | M. Pt. °C | Mass. Spec. m/e |
|-------|-----------------|---------------------|----|----|-------------------------|-----------------------|-----------------|-------------|-----------------|
| 2-1 | CH ₂ | 4-methyl-phenyl | CH | CH | 2-(methylsulfonyl)ethyl | 4-methoxy-benzylamino | 10 | 85.7-86.6 | 502 |
| 2-2 | CH ₂ | 4-methyl-phenyl | CH | CH | 2-(methylsulfonyl)ethyl | NH ₂ | 10 | 169.1-170.0 | 385 |
| 2-3 | CH ₂ | 4-fluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | NH ₂ | 10 | | 386 |
| 2-4 | CH ₂ | 2,4-difluoro-phenyl | CH | CH | 2-(methylsulfonyl)ethyl | NH ₂ | 10 | 152.9-153.2 | 404 |
| 2-5 | CH ₂ | 4-fluoro-phenyl | CH | CH | 2-(methylthio)ethyl | NH ₂ | 14 | | 355 |
| 2-6 | CH ₂ | 4-ethoxyphenyl | CH | CH | 2-(methylsulfonyl)ethyl | NH ₂ | 14 | | 413 |
| 2-7 | CH ₂ | 2-fluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | NH ₂ | 14 | | 387 |
| 2-8 | CH ₂ | 2,6-difluorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | NH ₂ | 14 | | 405 |
| 2-9 | CH ₂ | 2-methoxyphenyl | CH | CH | 2-(methylsulfonyl)ethyl | NH ₂ | 14 | | 399 |
| 2-10 | CH ₂ | 2-chlorophenyl | CH | CH | 2-(methylsulfonyl)ethyl | NH ₂ | 14 | | 403 |

Preferred Embodiments

While the broadest definition of this invention is set forth in the Summary of the Invention, certain compounds of Formula (I) are preferred.

In certain preferred embodiments, R³ is hydrogen.

5 In certain preferred embodiments, A is -CH₂-, CH₂-CH₂- or -CH(CH₃)-; preferably -CH₂-.

In certain preferred embodiments, X is CH and Y is CH.

In other preferred embodiments, X is N and Y is CH.

In other preferred embodiments, X is CH and Y is N.

10 In certain preferred embodiments, B is aryl, preferably optionally substituted phenyl.

In other preferred embodiments, B is heteroaryl, preferably furyl, imidazolyl, pyridyl, thienyl, thiazolyl, benzothiazolyl or pyridazinyl.

In certain preferred embodiments R¹ is alkyl, cycloalkyl, cycloalkyl-alkyl, heterocyclyl, heterocyclylalkyl or heteroalkyl; more preferably heteroalkyl, especially alkylsulfonyl-alkyl
15 (e.g. 2-methylsulfonyl-ethyl).

In certain preferred embodiments R² is alkyl, more preferably methyl.

In other preferred embodiments R² is NH₂.

A particularly preferred group is (II) where X and Y are CH.

Within this group, in one preferred embodiment, B is phenyl optionally substituted from the group consisting of halo, alkoxy, and cyano, especially mono-substituted with fluoro
20 (e.g. 4-fluorophenyl); R¹ is alkylsulfonyl-ethyl, particularly 2-methylsulfonyl-ethyl; and R² is alkyl, particularly methyl or NH₂.

In another preferred group within (II), B is heteroaryl, preferably furyl, imidazolyl, pyridyl, thienyl, thiazolyl, benzothiazolyl or pyridazinyl; R¹ is alkylsulfonyl-ethyl, particularly 2-
25 methylsulfonyl-ethyl; and R² is alkyl, particularly methyl, or NH₂.

A number of different substituent preferences have been given above and following any of these substituent preferences results in a compound of the invention that is more preferred than one in which the particular substituent preference is not followed. However, these
30 substituent preferences are generally independent, although some preferences are mutually

exclusive, and following more than one of these preferences may result in a more preferred compound than one in which fewer of the substituent preferences are followed.

General Synthetic Scheme

5 Compounds of this invention can be made by the methods depicted in the reaction schemes shown below.

 The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Co., (Milwaukee, WI), Bachem (Torrance, CA), or Sigma (St. Louis, MO), Lancaster Synthesis (Pelham, N.C.), Maybridge Chemical Co.
10 LTD (Cornwall, United Kingdom) or are prepared by methods known to those skilled in the art following procedures set forth in references such as *Fieser and Fieser's Reagents for Organic Synthesis*, Volumes 1-17 (John Wiley and Sons, 1991); *Rodd's Chemistry of Carbon Compounds*, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989); *Organic Reactions*, Volumes 1-40 (John Wiley and Sons, 1991), *March's Advanced Organic Chemistry*, (John Wiley
15 and Sons, 4th Edition) and *Larock's Comprehensive Organic Transformations* (VCH Publishers Inc., 1989). These schemes are merely illustrative of some methods by which the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to one skilled in the art having referred to this disclosure.

 The starting materials and the intermediates of the reaction may be isolated and purified if
20 desired using conventional techniques, including but not limited to filtration, distillation, crystallization, chromatography and the like. Such materials may be characterized using conventional means, including physical constants and spectral data.


 Unless specified to the contrary, the reactions described herein take place at atmospheric pressure over a temperature range from about -78 °C to about 150 °C, more preferably from about
25 0 °C to about 125 °C and most preferably at about room (or ambient) temperature, e.g., about 20 °C.

Preparation of Compounds of Formula (I)

Schemes A-F describe methods to generate the compounds of Formula (I). One of skill in the art will recognize that groups R^1 , R^2 , R^3 , A and B may be present in protected form at any point in the Schemes and will be removed at the appropriate juncture.

- 5 Scheme A describes the synthesis of a compound of formula (I) wherein X and Y are CH and R^1 , R^2 , R^3 , A, and B are as defined in the Summary of the Invention.

Scheme A



In Step 1, reaction of a phenylsulfonyl compound of formula **1** (where Z is an appropriate leaving group such as fluoro or bromo) with a monosubstituted amine of formula **2** (wherein R^1 is as defined in the Summary of the Invention or a protected precursor thereof) gives a 4-aminophenylsulfonyl compound of formula **3**. The reaction is carried out at elevated temperature,

preferably in the range of 50 - 80° C and in the presence of a base such as potassium carbonate, triethylamine, and the like. Suitable solvents for the reaction are polar aprotic solvents such as DMF, DMSO, HMPA, and the like. In general, the compounds of formula 1 are commercially available or can be readily synthesized by those of ordinary skill in the art.

5 Compounds of formula (I) can be prepared from a compound of formula 3 by any of the following three steps: Step 2a, 2b, or 2c.

As shown in Step 2a, a compound of formula (I) can be prepared from a compound of formula 3 by reductive amination. Reaction of compound 3 with an aldehyde of formula 4a (wherein B is as defined in the Summary of the Invention or a protected precursor thereof) and a
10 suitable reducing agent (e.g. a hydride reducing agent such as NaBH(OAc)₃) gives a compound of Formula (I) (wherein A is CH₂). Suitable solvents for the reaction are halogenated hydrocarbons, such as dichloromethane, dichloroethane, and the like. See, for example, Example 3.

Alternatively, as shown in Step 2b, a compound of formula (I) can be prepared from a compound of formula 3 by direct nucleophilic alkylation. Reaction of compound 3 with a
15 compound of formula 4b (where B and A are defined in the Summary of the Invention or a protected precursor thereof and Z is an appropriate leaving group such as bromo and chloro) gives a compound of formula (I). The reaction is carried out in the presence of a base such as sodium hydride. Suitable solvents for the reaction are polar aprotic solvents such as DMF, DMSO, HMPA, and the like. This reaction is carried out at approximately room temperature to 70°C.
20 See, for example, Examples 2 and 4.


Alternatively, as shown in Step 2c, a compound of formula (I) can be prepared from a compound of formula 3 by acylation/reduction. Compound 3 is reacted with an acid chloride or carboxylic acid of formula 4c (where Z is a leaving group, such as chloro or -OH and B is as defined in the Summary of the Invention or a protected precursor thereof). If the compound of
25 formula 4c is a carboxylic acid a coupling agent, such as DCC, must also be present. This acylation is followed by reduction using a suitable reducing agent (typically a hydride reducing agent such as LAH, B₂H₆, BH₃DMS, and the like) to provide a compound of formula (I) (wherein A is CH₂). Suitable solvents for the reaction are polar, anhydrous solvents such as THF, ether, and the like.

Additional steps may be added to this general Scheme A as necessary to provide the desired compound of formula (I). As an example of an additional step, it may be necessary to protect functional groups in R¹ of compound 3 before preparing a compound of formula (I) by Step 2a, 2b, or 2c and then subsequently deprotect the functional group. For example, amine
5 functionality in a compound of formula 3, may be protected by treatment of 3 with di-*tert*-butyl dicarbonate, followed by removal of the *tert*-butoxy carbonyl protecting group after Step 2. Examples of protecting groups and their synthetic use can be found in T.W. Greene and P.G. Futs, Protective Groups in Organic Chemistry, (Wiley, 2nd ed. 1991) and Harrison and
10 Harrison *et al.*, Compendium of Synthetic Organic Methods, Vols. 1-8 (John Wiley and Sons. 1971-1996). These protection/deprotection steps are well known by those of ordinary skill in the art and need not be further elaborated.

Compounds of formula (I) may also be obtained by further modification of a functional group on a compound of formula (I). For example, compounds of formula (I), where R¹ is alkylsulfonylalkyl (e.g. methylsulfonyl ethyl) may be obtained by oxidation of the corresponding
15 alkylthioalkyl compound which in turn may be prepared via step 1 by treating a compound of formula 1 with the corresponding alkylthioalkyl amine.

Scheme B describes an alternative synthesis of a compound of formula (I) by sequential alkylation of an aromatic amine, 5, wherein X, Y, R¹, R², R³, A and B are as defined in the Summary of the Invention or protected precursors thereof, except that R² is not NR¹³R¹⁴

Scheme B



In Step 1, reaction of an aromatic amine of formula 5 with a nucleophilic alkylating agent of formula 6 (where B and A are as defined in the Summary of the Invention or a protected precursor thereof and Z is an appropriate leaving group such as bromo, and chloro) in the presence of a base (such as triethylamine (TEA) or diisopropylethyl amine) provides a compound of formula 7. Suitable solvents for this reaction are dichloromethane, THF, and the like.


In Step 2, a compound of formula 7 is reacted with an alkylating agent of formula 8 (wherein R¹ is as defined in the Summary of the Invention and Z is an appropriate leaving group such as bromo and chloro) to provide a compound of formula 9. The reaction is carried out in the presence of a base such as sodium hydride. Suitable solvents for the reaction are polar, aprotic solvents such as DMF, DMSO, HMPA, and the like.

In Step 3, oxidation of a compound of formula 9 with a suitable oxidizing agent, such as potassium peroxymonosulfate (OXONE™), MCPBA, and the like, provides a compound of formula (I). Suitable solvents for the reaction are alcohols, such as methanol and ethanol. See, for example, Example 1.

Scheme C describes the synthesis of a compound of formula (I) wherein X, Y, R², R³ and B are as defined in the Summary of the Invention (except that R² is not NR¹³R¹⁴), A is -CH₂- and R¹ is alkylsulfonylalkyl.

5

Scheme C




In Step 1 of Scheme C, reaction of an aromatic amine sulfide of formula 10 with an aldehyde of formula 11 and a suitable reducing agent (such as NaBH(OAc)₃) gives an amino substituted aromatic sulfide of formula 12.

In Step 2, an aromatic sulfide of formula 12 is reacted with a vinyl sulfone of formula 13 in the presence of a base such as sodium hydride to provide a sulfide of formula 14. Suitable solvents for the reaction are polar aprotic solvents such as DMF, DMSO, HMPA, and the like

In Step 3, oxidation of a sulfide of formula 14 with a suitable oxidizing agent, such as potassium peroxymonosulfate (OXONE TM), MCPBA, and the like, provides a compound of formula (I). Suitable solvents for the reaction are alcohols, such as methanol and ethanol. See, for example, Example 5.

5 4-anilino sulfides of formula 10 where X and Y are both CH are available from commercial suppliers such as Aldrich Chemical Co. Amino-pyridyl sulfides and amino-pyridazinyl sulfides of formula 10, where either of X and Y are N may be prepared from the corresponding amino-pyridines and amino-pyridazines by halogenation and alkylation with a thiolate as shown in Scheme C1.


Scheme C1



In Step 1, the heteroaromatic amine is treated with I₂ in DMSO as described in *Heterocycles* 1984, 1195 to give an iodinated product that is treated with NaSR₂ in DMF in Step 2 to displace the iodine to form a compound of formula 10 which is carried forward into Scheme C.

Alternatively, compounds of formula 10 where either of X and Y are N may also be prepared by thiolation of the corresponding halo nitro heteroaromatic compounds followed by reduction of the nitro group to an amine as shown in Scheme C2.


Scheme C2



5 In Step 1, a bromo-nitro-heteroaromatic compound is treated with NaSR_2 in DMF to displace the bromine to form the corresponding heteroaromatic sulfide. The sulfide is treated with TiCl_3 in acetone and NH_4OAc as described in *Chem. Soc. Perkin. Trans. I.* 1990, 673 to give a compound of formula 10 that is carried forward into Scheme C.

10 Intermediates of formula 12 from Scheme C may also be alkylated with an alkylating agent, $\text{R}^1\text{-Z}$, or reductively aminated with an aldehyde, RCHO , as shown in Scheme D to furnish corresponding compounds of Formula I after oxidation.


Scheme D



Scheme E describes an alternative synthesis of a compound of formula (I) wherein X and Y are CH, R² is alkylsulfonylalkyl and R¹, R³, A, and B are as defined in the Summary of the Invention, wherein the alkylsulfonylalkyl group is introduced before the introduction of the A-B group.

5

Scheme E



10 In Step 1, reaction of a compound of formula 5 with a vinyl sulfone of formula 13 in the presence of a base such as sodium hydride provides a compound of formula 15. Suitable solvents for the reaction are polar aprotic solvents such as DMF, DMSO, HMPA, and the like.


In Step 2, oxidation of a compound of formula 15 with a suitable oxidizing agent, such as such as potassium peroxymonosulfate (OXONE[™]), MCPBA, and the like, provides a sulfone of formula 16. Suitable solvents for the reaction are alcohols, such as methanol and ethanol.

Compounds of formula 16 may be converted to compounds of formula (I) by following Scheme A: Steps 2a, 2b, or 2c above.

Compounds where R² is NH₂ may be prepared by the sequence shown in Scheme F.

5

Scheme F




10 In step i) a compound of formula 5 is acylated with an acid chloride of formula 17 to give an amide that is chlorosulfonylated and aminated with bis(p-methoxybenzyl)amine (PMB) steps ii) and iii) to give a benzene sulfonamide of formula 18.

15 In step iv), reduction of the amide in 18 gives an amine which is then subsequently elaborated in steps v) and vi) by alkylation with a vinyl sulfone to give a compound of formula (I) where R¹ is 2-alkylsulfonyl-ethyl, or by alkylation with R¹-Z or by reductive amination with an aldehyde RCHO as described earlier.

Additional compounds where R^2 is $NR^{13}R^{14}$ or NH_2 or may be prepared by Schemes G and H respectively as shown below.


Scheme G



Acylation of a para-methylthioethylamino sulfonamide 19 in the absence of base with an acylating agent $BC(O)Cl$ in an inert solvent provides the monoacylated intermediate 20.

Subsequent acylation of the sulfonamido group of 20 provides the bisacylated product 21, which is then reduced to give intermediate 22. Oxidation then provides 23, i.e., compounds Formula I where $A = CH_2$ and $R^2 = NHR^{13}$ (R^{13} is alkyl).

Scheme H



Alternatively, as shown in Scheme H, the amide group in monoacylated intermediate **20** may be reduced to provide a compound of Formula **24**, i.e., compounds of Formula I where R^2 is NH_2 . Compounds of Formula **24** may be further elaborated by oxidation to provide compounds of Formula **25** where R^1 is alkylsulfonyl-alkyl group.

In light of this disclosure, a person of ordinary skill in the art may readily prepare any desired compound of formula (I) by following the preceding Schemes.

General Utility

The compounds of the invention are inhibitors of prostaglandin G/H Synthase I and II (COX I and COX II), especially COX II, *in vitro*, and as such are expected to possess both anti-inflammatory and analgesic properties *in vivo*. See, for example, Goodman and Gilman's "The

Pharmacological Basis of Therapeutics”, Ninth Edition, McGraw Hill, New York, 1996, Chapter 27. The compounds, and compositions containing them, are therefore useful as anti-inflammatory and analgesic agents in mammals, especially humans. They find utility in the treatment of fever, inflammation and pain caused by conditions such as rheumatic fever, symptoms associated with influenza or other viral infections, low back and neck pain, dysmenorrhoea, headache, toothache, sprains, myositis, synovitis, arthritis (rheumatoid arthritis and osteoarthritis), gout, ankylosing spondylitis, bursitis, burns or injuries. They may be used to inhibit prostanoid-induced smooth muscle contractions (e.g., in the treatment of dysmenorrhoea, premature labor and asthma) and to treat autoimmune disorders (such as systemic lupus erythematosus and type I diabetes).

As inhibitors of prostaglandin G/H Synthase, the compounds of this invention are also expected to be useful in the prevention and treatment of cancer, in particular colon cancer. It has been shown that COX-2 gene expression is upregulated in human colorectal cancers and that drugs that inhibit prostaglandin G/H Synthase are effective in animal models of cancer (Eberhart, C.E., *et. al.*, *Gastroenterology*, **107**, 1183-1188, (1994), and Ara, G. and Teicher, B.A., *Prostaglandins, Leukotrienes and Essential Fatty Acids*, **54**, 3-16, (1996)). In addition, there is epidemiological evidence that shows a correlation between use of drugs that inhibit prostaglandin G/H synthase and a reduced risk of developing colorectal cancer, (Heath, C.W. Jr., *et. al.*, *Cancer*, **74**, No. 10, 2885-8, (1994)).

The compounds of this invention are also expected to be useful in the prevention and treatment of Alzheimer’s disease. Indomethacin, an inhibitor of prostaglandin G/H synthase, has been shown to inhibit the cognitive decline of Alzheimer’s patients, (Rogers, J., *et. al.*, *Neurology*, **43**, 1609, (1993)). Also, the use of drugs which inhibit prostaglandin G/H synthase has been linked epidemiologically with a delayed onset of Alzheimer’s disease, (Breitner, J.C.S., *et. al.*, *Neurobiology of Aging*, **16**, No. 4, 523, (1995) and *Neurology*, **44**, 2073, (1994)).

Testing

The anti-inflammatory activity of the compounds of this invention may be assayed by measuring the ability of the compound to inhibit COX I and COX II, especially COX II, *in vitro*, using a radiometric assay, as described in more detail in Example 9. It may also be

assayed by *in vivo* assays such as the Rat Carrageenan Paw and Rat Air-Pouch assays, as described in more detail in Examples 10 and 11. The analgesic activity of the compounds of this invention may be assayed by *in vivo* assays such as the Randall-Selitto assay and the rat arthritis pain model, as described in Example 12.

5

Administration and Pharmaceutical Composition

In general, the compounds of this invention will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. The actual amount of the compound of this invention, i.e., the active ingredient, will depend upon numerous factors such as the severity of the disease to be treated, the age and relative health of the subject, the potency of the compound used, the route and form of administration, and other factors.

10

Therapeutically effective amounts of compounds of formula (I) may range from approximately 0.05-35 mg per kilogram body weight of the recipient per day, preferably about 0.15-7 mg/kg/day, most preferably about 0.35 mg/kg/day to 3 mg/kg/day. Thus, for administration to a 70 kg person, the dosage range would preferably be about 10.5 mg to 500 mg per day, most preferably about 25 mg to 200 mg per day.

15

In light of this disclosure, a person of ordinary skill in the art will have no difficulty in determining what a therapeutically effective amount is.

20

In general, compounds of this invention will be administered as pharmaceutical compositions by any one of the following routes: oral, systemic (e.g., transdermal, intranasal or by suppository), or parenteral (e.g., intramuscular, intravenous or subcutaneous) administration. The preferred manner of administration is oral using a convenient daily dosage regimen which can be adjusted according to the degree of affliction. Compositions can take the form of tablets, pills, capsules, semisolids, powders, sustained release formulations, solutions, suspensions, elixirs, aerosols, or any other appropriate compositions.

25

The choice of formulation depends on various factors such as the mode of drug administration (e.g., for oral administration, formulations in the form of tablets, pills or capsules are preferred) and the bioavailability of the drug substance. Recently, pharmaceutical formulations have been developed especially for drugs that show poor bioavailability based

30

upon the principle that bioavailability can be increased by increasing the surface area i.e., decreasing particle size. For example, U.S. Pat. No. 4,107,288 describes a pharmaceutical formulation having particles in the size range from 10 to 1,000 nm in which the active material is supported on a crosslinked matrix of macromolecules. U.S. Pat. No. 5,145,684 describes the production of a pharmaceutical formulation in which the drug substance is pulverized to nanoparticles (average particle size of 400 nm) in the presence of a surface modifier and then dispersed in a liquid medium to give a pharmaceutical formulation that exhibits remarkably high bioavailability.

The compositions are comprised of, in general, a compound of formula (I) in combination with at least one pharmaceutically acceptable excipient. Acceptable excipients are non-toxic, aid administration, and do not adversely affect the therapeutic benefit of the compound of formula (I). Such excipient may be any solid, liquid, semi-solid or, in the case of an aerosol composition, gaseous excipient that is generally available to one of skill in the art .

Solid pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk and the like. Liquid and semisolid excipients may be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, etc. Preferred liquid carriers, particularly for injectable solutions, include water, saline, aqueous dextrose, and glycols.

Compressed gases may be used to disperse a compound of this invention in aerosol form. Inert gases suitable for this purpose are nitrogen, carbon dioxide, etc.

Other suitable pharmaceutical excipients and their formulations are described in *Remington's Pharmaceutical Sciences*, edited by E. W. Martin (Mack Publishing Company, 18th ed., 1990).

The level of the compound in a formulation can vary within the full range employed by those skilled in the art. Typically, the formulation will contain, on a weight percent (wt%) basis, from about 0.01-99.99 wt% of a compound of formula (I) based on the total formulation, with the balance being one or more suitable pharmaceutical excipients. Preferably, the

compound is present at a level of about 1-80 wt%. Representative pharmaceutical formulations containing a compound of formula (I) are described in Example 8.

A person of ordinary skill in the art will have no difficulty, having regard to that skill and this disclosure in determining how to make a suitable formulation.

5

EXAMPLES

The following preparations and examples are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.


10

Abbreviations used in the examples are defined as follows: "HCl" for hydrochloric acid, "DMF" for dimethylformamide, "NaOH" for sodium hydroxide, "DMSO" for dimethylsulfoxide, "THF" for tetrahydrofuran, "BINAP" for 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl.

15

Example 1

Synthesis of 4-[N,N-(benzyl)(4-fluorobenzyl)amino]phenyl methyl sulfone (1-42)



Step 1

To 5.0 mL (40.19 mmol) of 4-(methylthio)aniline and 8.4 mL (60.28 mmol) triethylamine dissolved in 30 mL dichloromethane was added 5.0 mL (40.19 mmol) 4-fluorobenzyl bromide. The mixture was stirred at room temperature for 12 h, partitioned between dichloromethane and sat. aqueous ammonium chloride, dried over MgSO₄ and concentrated. Purification by column chromatography, eluting with ethyl acetate/hexane,

20

provided 3.10 g of 4-[(4-fluorobenzyl)amino]thioanisole, along with recovered starting material.

Step 2


100 mg (0.40 mmol) of 4-[(4-fluorobenzyl)amino]thioanisole was dissolved in 2 mL of DMF, to which 76 mg (0.44 mmol) of benzyl bromide was added, followed by 19 mg (0.80 mmol) of NaH. The mixture was warmed to 50°C and stirred 48 h, quenched with water and concentrated. The residue was partitioned between ethyl acetate and water, dried over MgSO₄ and concentrated to provide 70 mg (0.207mmol) of crude 4-[N,N-(benzyl)(4-fluorobenzyl)amino]thioanisole.

Step 3

The crude 4-[N,N-(benzyl)(4-fluorobenzyl)amino]thioanisole was dissolved in 1.5 mL of methanol, to which 254 mg (0.414 mmol) of OXONE™ was added. The reaction was stirred at room temperature for 12 h, partitioned between ethyl acetate and water, dried over MgSO₄ and concentrated to provide 36 mg of 4[N,N-(benzyl)(4-fluorobenzyl)amino]phenyl methyl sulfone.

Example 2

Synthesis of 4-[N,N-(pyrrolidin-3-yl)(4-fluorobenzyl)amino]phenyl methyl sulfone (1-40)



Step 1

To 500 mg (2.87 mmol) 4-fluorophenyl methyl sulfone dissolved in 4 mL DMF was added 247mg (2.87mmol) 3-aminopyrrolidine followed by 793 mg (5.74 mmol) potassium carbonate. The mixture was heated to 70°C and stirred for 48 h. Upon cooling to room

temperature, the mixture was partitioned between EtOAc and water, dried over MgSO₄ and concentrated to provide 708 mg of 4-(pyrrolidin-3-ylamino)phenyl methyl sulfone, pure by ¹H NMR.

Step 2

4-(pyrrolidin-3-ylamino)phenyl methyl sulfone (2.96 mmol) was dissolved in 5 mL of THF, to which 645 mg (2.96 mmol) di-*tert* butyl dicarbonate was added. After 1 h, the mixture was partitioned between EtOAc and water, dried over MgSO₄, concentrated and the product crystallized from dichloromethane/hexane to provide 461 mg of 4-(N-BOC-pyrrolidin-3-ylamino)phenyl methyl sulfone, pure by ¹H NMR.

Step 3

4-(N-BOC-pyrrolidin-3-ylamino)phenyl methyl sulfone (1.35 mmol) was dissolved in 4 mL of DMF, to which 168 μ L (1.35 mmol) of 4-fluorobenzyl bromide was added, followed by 62 mg (2.70 mmol) of NaH. The mixture was warmed to 70°C and stirred 48 h, then quenched with water, partitioned between ethyl acetate and water, dried over MgSO₄ and concentrated.

The product was crystallized from dichloromethane/hexane to provide 380 mg of 4-[N,N-(N-BOC-pyrrolidin-3-yl)(4-fluorobenzyl)amino]phenyl methyl sulfone, pure by ¹H NMR; mp 174.4-178.0°C. Anal. Calcd. For C₂₃H₂₉FN₂O₄S·0.5H₂O C, 60.37; H, 6.61; N, 6.12. Found C, 60.61; H, 6.40; N, 6.34. (38)


To 200 mg (0.45 mmol) of 4-[N,N-(N-BOC-pyrrolidin-3-yl)(4-fluorobenzyl)amino]phenyl methyl sulfone dissolved in 5 mL of dichloromethane was added 2 mL of trifluoroacetic acid. The reaction was stirred at room temperature for 3 h, partitioned between dichloromethane and aqueous saturated NaHCO₃, was dried over MgSO₄ and concentrated to provide 148 mg of 4-[N,N-(pyrrolidin-3-yl)(4-fluorobenzyl)amino]phenyl methyl sulfone, pure by ¹H NMR, mp 124.0-124.3°C. Anal. Calcd. For C₁₈H₂₁FN₂O₂S C, 62.05; H, 6.07; N, 8.04. Found C, 61.29; H, 6.00; N, 7.92. (40)

Following the procedure of Example 1, but replacing 3-aminopyrrolidine in Step 1 with the appropriate amine gave the compounds #24-30, 32, 36-37 and 39-41 of Table 1.

Following the procedure of Example 1, but replacing 3-aminopyrrolidine in Step 1 with n-butylamine and replacing 4-fluorobenzyl bromide in Step 3 with the appropriate aralkyl bromide gave the compounds # 47-54 of Table 1.

Example 3

Synthesis of 4-[N,N-(butyl)(thiophen-2-ylmethyl)amino]phenyl methyl sulfone (1-22)



Step 1


To 1.0 g (5.74 mmol) 4-fluorophenyl methyl sulfone dissolved in 5 mL DMF was added 840 μ L (11.48 mmol) butylamine followed by 873 mg (6.31 mmol) potassium carbonate. The mixture was heated to 60°C and stirred for 48 h. Upon cooling to room temperature, the mixture was partitioned between EtOAc and water, dried over MgSO_4 and concentrated. Purification by column chromatography, eluting with dichloromethane/hexane, provided 600 mg product 4-(butylamino)phenyl methyl sulfone, pure by ^1H NMR.

Step 2

To 250 mg (1.1 mmol) of 4-(butylamino)phenyl methyl sulfone and 103 μ L 2-thiophenecarboxaldehyde dissolved in 5 mL dichloromethane was added 350 mg (1.65 mmol) sodium triacetoxyborohydride, followed by 50 μ L of acetic acid. The reaction mixture was stirred at room temperature for 12 h. The mixture was then partitioned between ethyl acetate and brine, dried over MgSO_4 and concentrated. Purification by HPLC chromatography provided 58 mg of 4-[N,N-(butyl)(thiophen-2-ylmethyl)amino]phenyl methyl sulfone (22).

Example 4

Synthesis of 4-{N,N-[2-(methylsulfonyl)ethyl](4-fluorobenzyl)amino}phenyl methyl sulfone
(1-35)



Step 1

To 8.28 g (47.52 mmol) of 4-fluorophenyl methyl sulfone dissolved in 20 mL DMF was added 5.20 g (57.03 mmol) of 2-(methylthio)ethyl amine (1.2 eq.), followed by 13.13 g (95.04 mmol, 2 eq.) potassium carbonate. The mixture was heated to 65°C and stirred for 12 h. Upon cooling to room temperature, the mixture was partitioned between EtOAc and brine, dried over MgSO₄ and concentrated. Column chromatography, eluting with ethyl acetate/hexane, provided 5.53 g of 4-[2-(methylthio)ethylamino]phenyl methyl sulfone, pure by ¹H NMR.

Step 2

To 2.5 g of 4-[2-(methylthio)ethylamino]phenyl methyl sulfone (10.19 mmol) dissolved in 10 mL DMF was added 1.26 mL (10.19 mmol) of 4-fluorobenzyl bromide and 468 mg (20.38 mmol) of NaH. After stirring 1 h at room temperature, the reaction was quenched with water and partitioned between EtOAc and water, dried over MgSO₄ and concentrated. Purification by column chromatography, eluting with ethyl acetate/hexane, provided 2.44 g product 4-[N,N-(methylthioethyl)(4-fluorobenzyl)amino]phenyl methyl sulfone, pure by ¹H NMR.

Step 3

To 2.44 g (6.91 mmol) of 4-[N,N-[2-(methylthio)ethyl](4-fluorobenzyl)amino]phenyl methyl sulfone dissolved in 40 mL MeOH was added 8.5 g (13.83 mmol) OXONE™, followed by slow addition of 5 mL H₂O. After stirring at room temperature for 7 h, the mixture was


partitioned between EtOAc and water, dried over MgSO_4 and concentrated. The product was crystallized from CH_2Cl_2 to obtain 600 mg of 4-{N,N-[2-(methylsulfonyl)ethyl](4-fluorobenzyl)amino}phenyl methyl sulfone, pure by ^1H NMR, mp 168.6-172.7 °C.

Following the procedure of Example 4, but replacing 2-(methylthio)ethyl amine in step 1 with 2-(ethylthio)ethyl amine gave 4-{N,N-[2-(ethylsulfonyl)ethyl](4-fluorobenzyl)amino}phenyl methyl sulfone. Mp. 109.6-110.7 °C Anal. Calcd. For $\text{C}_{18}\text{H}_{22}\text{FNO}_4\text{S}_2$ C, 54.12; H, 5.55; N, 3.51. Found C, 53.72; H, 5.48; N, 3.58. (34)

Following the procedure of Example 4, but replacing 4-fluorobenzyl bromide in Step 3 with 2,4-difluorobenzyl bromide gave 4-{N,N-[2-(methylsulfonyl)ethyl](2,4-difluorobenzyl)amino}phenyl methyl sulfone. Mpt. 149.4-150.4 °C. Anal. Calcd. For $\text{C}_{17}\text{H}_{19}\text{F}_2\text{NO}_4\text{S}_2 \cdot 0.25\text{H}_2\text{O}$ C, 50.05; H, 4.82; N, 3.43. Found C, 50.04; H, 4.63; N, 3.45. (44)

Example 5

Synthesis of 4-{N,N-[2-(methylsulfonyl)ethyl](pyridin-2-ylmethyl)amino}phenyl methyl sulfone (1-45)



Step 1

To 2.0 mL (16.07 mmol) of 4-(methylthio)aniline dissolved in 25 mL dichloromethane was added 1.52 mL (16.07 mmol) 3-pyridinecarboxaldehyde, followed by 5.11 g (24.11 mmol) sodium triacetoxyborohydride. The mixture was stirred at room temperature for 4 h, partitioned between EtOAc and brine, dried over MgSO_4 and concentrated. Column chromatography, eluting with ethyl acetate/hexane, provided 3.70 g of 4-[(pyridin-2-ylmethyl)amino]thioanisole, pure by ^1H NMR.

Step 2


To 500 mg (2.17 mmol) of 4-[(pyridin-2-ylmethyl)amino]thioanisole, dissolved in 10 mL *N,N*-dimethylformamide, was added 230 mg (2.17 mmol) methyl vinyl sulfone followed by 50 mg (2.17 mmol) sodium hydride. The mixture was stirred at room temperature for 0.5 h, partitioned between EtOAc and brine, dried over MgSO₄ and concentrated. Crude 4-{N,N-[2-(methylsulfonyl)ethyl](pyridin-2-ylmethyl)amino}thioanisole was obtained in 98% yield (730 mg) and was pure by ¹H NMR.

Step 3

To 718 mg (2.13 mmol) of 4-{N,N-[2-(methylsulfonyl)ethyl](pyridin-2-ylmethyl)amino}thioanisole dissolved in 10 mL methanol was added 2.62 g (4.27 mmol) oxone followed by 500 µL water. The mixture was stirred at room temperature for 1 h, then partitioned between EtOAc and water, adding 1 N NaOH until the aqueous phase was neutral. The organic layer was then dried over MgSO₄ and concentrated to give 4-{N,N-[2-(methylsulfonyl)ethyl](pyridin-2-ylmethyl)amino}phenyl methyl sulfone in 57% yield (446 mg), pure by ¹H NMR.

Example 6

Synthesis of 4-{N,N-[2-(methylsulfonyl)ethyl](4-bromobenzyl)amino}phenyl methyl sulfone (1-12)



Step 1

To 8.94ml (1.83mmol) of 4-(methylthio)aniline dissolved in 100ml DMF was added 1.72g (71.67mmol) NaH at 0°C, followed with 6.29 ml (71.80 mmol) methyl vinyl sulfone. The mixture was stirred at room temperature for 14h, quenched with MeOH and concentrated.

The residue was dissolved in CH₂Cl₂, washed with aqueous HCl (1M) (100mlx2), then water, dried over MgSO₄ and concentrated. The crude product was purified by flash chromatography (Hexane : ethyl acetate 1:1) to provide 4-[2-methylsulfonyl)ethylamino]phenyl methylthiol (3.60 g) as a yellow solid.

5 Step 2


A solution of 4-[2-methylsulfonyl)ethylamino]phenyl methylthiol (3.60g; 14.67 mmol) in 200ml MeOH and 50ml THF was cooled to 0°C, to which a mixture of 13.57g (22.07mmol) OXONE™ with 50ml warm water was added. The mixture was stirred at room temperature for 30 minute. The OXONE™ solid was filtered off and the filtrate was concentrated. A brown solid precipitated out during the concentrated, which was filtered out to provide 4-[2-methylsulfonyl)ethylamino]phenyl methyl sulfone (2.38g).

Step 3

To a solution of 4-[2-methylsulfonyl)ethylamino]phenyl methyl sulfone (0.80g; 2.88mmol) in 20ml DMF was added 0.104g (4.33mmol) NaH followed by 1.25 ml (9.21mmol) 4-bromobenzyl bromide. The mixture was stirred at room temperature for 1h, quenched with MeOH and concentrated. The residue was purified by prep-TLC (hexane: ethyl acetate 1:2) to provide 0.765g product 4-{N,N-[2-(methylsulfonyl)ethyl](4-bromobenzyl)amino }phenyl methyl sulfone as a white foam.

Example 7

Synthesis of 4-{N,N-[2-(methylsulfonyl)ethyl](4-ethoxybenzyl)amino }phenyl methyl sulfone (1-65)



Step 1

To 8.94ml (1.83mmol) of 4-(methylthio)aniline dissolved in 100ml DMF was added 1.72g (71.67mmol) NaH at 0°C, followed with 6.29 ml (71.80 mmol) methyl vinyl sulfone.

The mixture was stirred at room temperature for 14h, quenched with MeOH and concentrated.

- 5 The residue was dissolved in CH₂Cl₂, washed with aqueous HCl (1M) (100mlx2), then water, dried over MgSO₄ and concentrated. The crude product was purified by flash chromatography (Hexane : ethyl acetate 1:1) to provide 4-[2-methylsulfonyl)ethylamino]phenyl methylthiol (3.60 g) as a yellow solid.

Step 2

- 10 A solution of 4-[2-methylsulfonyl)ethylamino]phenyl methylthiol (3.60g; 14.67 mmol) in 200ml MeOH and 50ml THF was cooled to 0°C, to which a mixture of 13.57g (22.07mmol) OXONETM with 50ml warm water was added. The mixture was stirred at room temperature for 30 minute. The OXONETM solid was filtered off and the filtrate was concentrated. A brown solid precipitated out during the concentrated, which was filtered out to provide 4-[2-methylsulfonyl)ethylamino]phenyl methyl sulfone (2.38g).
- 15

Step 3


To a solution of 4-[2-methylsulfonyl)ethylamino]phenyl methyl sulfone (0.548g; 1.97mmol) and 0.73g (3.95 mmol) 4-ethoxybenzoyl chloride in anhydrous CH₂Cl₂ was added 0.32 ml (3.95mmol) pyridine. The mixture was warmed to 45 °C and stirred for 14h.

- 20 The reaction mixture was concentrated and the residue was purified with prep-TLC (hexane: ethyl acetate 1:3) to provide 4-{N,N-[2-(methylsulfonyl)ethyl](4-ethoxybenzyl)amino}phenyl methyl sulfone (0.664 g) as a white foam.

- The above product, 4-{N,N-[2-(methylsulfonyl)ethyl](4-ethoxybenzyl)amino}phenyl methyl sulfone (0.42g; 0.98 mmol), was dissolved in 20ml anhydrous toluene, to which
- 25 0.098ml (0.98mmol) BH₃•Me₂S (10.0 – 10.2 M) complex was added. The mixture was refluxed with stirring for 18 h, and quenched with aqueous NaHCO₃ (8ml). The toluene layer was separated and the aqueous layer was extracted with CH₂Cl₂ (100ml x3). The organic layers were combined, dried over MgSO₄ and concentrated. The residue was purified with prep-plate (hexane : ethyl acetate 1:2) to provide 4-{N,N-[2-(methylsulfonyl)ethyl](4-ethoxybenzyl)amino}phenyl methyl sulfone (0.143g) product as a white foam.
- 30

Example 8

Synthesis of 4-[N,N-(4-methylsulfonylphenyl)(4-fluorobenzyl)amino]phenyl methyl sulfone (1-68) and 4-[N,N-(4-methylthiophenyl)(4-fluorobenzyl)amino]phenyl methyl sulfone (1-69)



Step 1

To 150mg (0.64mmol) 4-bromo-phenyl methyl sulfone, 7.8mg (2%) tris(dibenzylideneacetone)dipalladium(0), 10.6mg (4%) BINAP, and 277mg (0.85mmol) Cs₂CO₃ in 2mL toluene was added 63.5 μ L (0.51mmol) 4-(methylthio)aniline. The mixture was heated to 100⁰C under N₂ and was stirred for 48h. The mixture was cooled, diluted with ether, filtered through celite, and concentrated. Purification by column chromatography, eluting with EtOAc/hexane, provided 128mg product 4-(4-methylsulfonyl-phenylamino)phenyl methyl sulfide, pure by ¹H NMR.

Step 2

To 191mg (0.65mmol) 4-(4-methylsulfonyl-phenylamino)phenyl methyl sulfide in 3mL DMF was added 27mg (0.68mmol) sodium hydride. The mixture was stirred under N₂ for 15min, then 122 μ L (0.98mmol) p-fluorobenzylbromide was added. The mixture was stirred for 18h. The mixture was partitioned between EtOAc and water, dried over Na₂SO₄ and concentrated. Purification by column chromatography, eluting with EtOAc/hexane, provided

118mg product 4-{N,N-(4-methylsulfonyl-phenyl)(4-fluorobenzyl)amino]phenyl methyl sulfide, pure by ¹H NMR and LCMS.


Step 3

5 To 138mg (0.34mmol) 4-{N,N-(4-methylsulfonyl-phenyl)(4-fluorobenzyl)amino]phenyl methyl sulfide and 423mg (0.69mmol) oxone in 3mL methanol was added 3drops (10% by vol) water. The mixture was stirred at room temperature for 18h. The mixture was partitioned between EtOAc and water, dried over Na₂SO₄ and concentrated. Purification by column chromatography, eluting with EtOAc/hexane, provided 115mg product 4-[N,N-(4-

10 methylsulfonyl-phenyl)(4-fluorobenzyl)amino]phenyl methyl sulfone, pure by ¹H NMR and LCMS.

Example 9

Synthesis of 4-{N,N-(3-oxo-butyl)(4-fluorobenzyl)amino]phenyl methyl sulfone (1-70)



Step 1

15 To 2.9mL (23mmol) 4-methylthio-aniline and 2.9mL (23mmol) 4-fluorobenzylbromide in 50mL CH₂Cl₂ was added 6.5mL (47mmol) triethylamine. The mixture was stirred at rt for 18h. The mixture was washed with water, dried over Na₂SO₄ and concentrated. Purification by

20 column chromatography, eluting with EtOAc/hexane, provided 2.2g product 4-(4-fluorobenzylamino)phenyl methyl sulfide, pure by ¹H NMR.

Step 2

25 To 100mg (0.40mmol) 4-(4-fluorobenzylamino)phenyl methyl sulfide in 1mL dioxane and 1mL phosphate buffer [1:4 KH₂PO₄/K₂HPO₄ with pH 7] was added dropwise 40μL

(0.48mmol) methyl vinyl ketone. The biphasic mixture was stirred at room temperature for 18h. Another 40 μ L (0.48mmol) methyl vinyl ketone was added, and the mixture was stirred for 18h. The mixture was extracted with ether, which was washed with water, dried over Na_2SO_4 and concentrated. Purification by column chromatography, eluting with


- 5 EtOAc/hexane, did not provide separation. The crude 55mg product 4-[N,N-(3-oxo-butyl)(4-fluorobenzyl)amino]phenyl methyl sulfide was used directly in Step 3.

Step 3

- To 55mg (0.17mmol) 4-[N,N-(3-oxo-butyl)(4-fluorobenzyl)amino]phenyl methyl sulfide in 2mL methanol was added 215mg (0.35mmol) oxone and 2 drops (10% by vol) water. The mixture was stirred at room temperature for 18h. The mixture was partitioned between EtOAc and water, dried over Na_2SO_4 and concentrated. Purification by preparative TLC, developing with EtOAc/hexane, provided 5.4mg 4-[N,N-(3-oxobutyl)(4-fluorobenzyl)amino]phenyl methyl sulfone, pure by ^1H NMR and LCMS.

Example 10

4-[(2-methylsulfonyl-ethyl)-(4-methyl-benzyl)-amino]-benzenesulfonamide (2-2) and
4-[(2-methylsulfonyl-ethyl)-(4-methyl-benzyl)amino]-N-(4-methoxy-benzyl)-
20 benzenesulfonamide (2-1)



Step 1

- To a 0 °C dichloromethane (300 mL) solution of 25 mL (274 mmol) of aniline was added 16 mL (123 mmol) para-toluoyl chloride over 10 min. The mixture was stirred at room temperature for 0.5 h, treated with 200 mL of ether and filtered immediately. The filtrate was

washed with 1 M HCl (2 x 50 mL), 0.1 M NaOH (2 x 50 mL), and saturated aqueous ammonium chloride, dried over Na₂SO₄ and concentrated. Para-methylbenzanilide (17.3 g) was obtained as a tan solid and used directly.

Steps ii) and iii)

- 5 Chlorosulfonic acid (5 mL) was cooled to 0 °C under a nitrogen atmosphere and treated with para-methylbenzanilide (850 mg, 4.0 mmol). The resulting solution was stirred at room temperature for 3 h, recooled with an ice-bath, treated with ca. 25 g of ice, ca. 100 mL of saturated sodium bicarbonate and bis(para-methoxybenzyl)amine (prepared according to *J. Org. Chem.* **1992**, 57, 7056, 1.1 g, 4.4 mmol) dissolved in ca. 50 mL of dichloromethane. The
- 10 biphasic mixture was stirred vigorously at room temperature for 16 h. The layers were separated and the aqueous phase was extracted with dichloromethane, washed with brine and dried over Na₂SO₄. Purification by column chromatography, eluting with 1:3 ethyl acetate/hexane, provided 1.4 g of 4-[(4-methyl-benzoyl)-amino]-[N,N-bis(4-methoxy-benzyl)]-benzenesulfonamide.

15 Step iv)

- 4-[(4-methyl-benzoyl)-amino]-[N,N-bis(4-methoxy-benzyl)]-benzenesulfonamide (1.4 g, 2.6 mmol) was dissolved in toluene (60 mL), treated with borane methyl sulfide complex (0.57 mL, 5.7 mmol) and heated to reflux for 2 h. Upon cooling, the mixture was quenched with Na₂SO₄(H₂O)₁₀, partitioned between pH 4 buffer and ethyl acetate, and dried over MgSO₄.
- 20 Purification by column chromatography, eluting with 2:3 ethyl acetate/hexane, provided 1.03 g of 4-[(4-methyl-benzyl)-amino]-[N,N-bis(4-methoxy-benzyl)]-benzenesulfonamide.

Step v)

- 4-[(4-methyl-benzyl)-amino]-[N,N-bis(4-methoxy-benzyl)]-benzenesulfonamide (1.03 g, 2.6 mmol) was dissolved in 6 mL of DMF at room temperature, to which methylvinyl
- 25 sulfone (0.175 mL, 2.0 mmol) and sodium hydride (95%, 60 mg, 2.4 mmol) were added. The reaction was stirred at room temperature for 1.5 h, partitioned between ethyl acetate and water, dried over MgSO₄ and purified by column chromatography, eluting with 1:4 acetone/hexane, to provide 935 mg of 4-[(2-methylsulfonyl-ethyl)-(4-methyl-benzyl)-amino]-[N,N-bis(4-methoxy-benzyl)]-benzenesulfonamide.

Step vi)


4-[(2-methylsulfonyl-ethyl)-(4-methyl-benzyl)-amino]-[N,N-bis(4-methoxybenzyl)]-benzenesulfonamide (730 mg, 1.17 mmol) was dissolved in dichloromethane (5 mL) at room temperature and treated with trifluoroacetic acid (5 mL). After 6 h, the volatiles were removed on a rotary evaporator and the residue was partitioned between aqueous sodium bicarbonate and ethyl acetate. Following drying over Na_2SO_4 and removal of the volatiles, the mixture was purified by column chromatography eluting with 2:3 ethyl acetate/hexane. The first product to elute was 4-[(2-methylsulfonyl-ethyl)-(4-methyl-benzyl)amino]-N-(4-methoxy-benzyl)-benzenesulfonamide (276 mg): mp 85.7-86.6 °C. The next product to elute was 4-[(2-methylsulfonyl-ethyl)-(4-methyl-benzyl)-amino]-benzenesulfonamide (184 mg): mp 169.1-170.0 °C, Anal. Calcd for $\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}_4\text{S}_2(\text{H}_2\text{O})_{0.6}$: C, 53.34; H, 5.77; N, 6.91. Found: C, 53.33; H, 5.74; N, 7.30.

Following the procedure of Example 10, but replacing p-toluoyl chloride in Step 1 with 4-fluorobenzoyl chloride in Step 1, 4-[(2-methylsulfonyl-ethyl)-(4-fluoro-benzyl)-amino]-benzenesulfonamide was produced as an amorphous glass: Anal. Calcd for $\text{C}_{17}\text{H}_{19}\text{FN}_2\text{O}_4\text{S}_2$: C, 49.73; H, 4.96; N, 7.25. Found: C, 49.39; H, 4.96; N, 6.86. (2-3)

Following the procedure of Example 10, but replacing p-toluoyl chloride in Step 1 with 2,4-difluorobenzoyl chloride, 4-[(2-methylsulfonyl-ethyl)-(2,4-difluorobenzyl)-amino]-benzenesulfonamide was produced: mp 152.9-153.2 °C. (2-4)

Example 11

3-[N,N-(2-methylsulfonyl-ethyl)(4-fluorobenzyl)amino]-pyridin-6-yl methyl sulfone (1-60)



To a solution of 2.0 g (9.85 mmol) 2-bromo-5-nitropyridine dissolved in 8 mL DMF was added 690 mg (9.85 mmol) of sodium thiomethoxide. The mixture was stirred at room temperature for 1 h, and partitioned between ethyl acetate and water, dried over MgSO_4 and concentrated to obtain 1.13 g of 2-methylthio-5-nitro-pyridine, pure by ^1H NMR.

5

To a separatory funnel containing 646 mg (3.796 mmol) 2-methylthio-5-nitro-pyridine dissolved in 20 mL acetone and 150 mL 4 M ammonium acetate was added 26.57 mL (26.57 mmol) of a 1M solution of TiCl_3 dissolved in $\text{CH}_2\text{Cl}_2/\text{THF}$. The mixture was shaken for 5 min, ethyl acetate added, and partitioned, dried over MgSO_4 and concentrated. Purification by column chromatography, eluting with ethyl acetate/hexane, provided 185 mg product, 5-amino-2-methylthio-pyridine, pure by ^1H NMR.

10

To 344 μL (3.20 mmol) of 4-fluorobenzaldehyde and 449 mg (3.20 mmol) of 5-amino-2-methylthio-pyridine dissolved in 8 mL dichloromethane was added 1.11 g (5.25 mmol) sodium triacetoxyborohydride. The reaction mixture was stirred at room temperature for 5 h, and partitioned between ethyl acetate and water, dried over MgSO_4 and concentrated to obtain 320 mg of 5-[(4-fluorobenzyl)amino]-pyridin-2-yl methyl sulfide, pure by ^1H NMR.

15

To 320 mg (1.29 mmol) of 5-[(4-fluorobenzyl)amino]-pyridin-2-yl methyl sulfide dissolved in 5 mL *N,N*-dimethylformamide was added 137 mg (1.29 mmol) methyl vinyl sulfone followed by 30mg (1.29 mmol) sodium hydride. The mixture was stirred at room temperature for 6 h, partitioned between EtOAc and brine, dried over MgSO_4 and concentrated. Crystallization from CH_2Cl_2 /hexane afforded 447 mg of product, 5-[*N,N*-(2-methylsulfonyl-ethyl)(4-fluorobenzyl)amino]-pyridin-2-yl methyl sulfide, pure by ^1H NMR.

20


25

To 447 mg (1.26 mmol) of 5-[*N,N*-(2-methylsulfonyl-ethyl)(4-fluorobenzyl)amino]-pyridin-2-yl methyl sulfide dissolved in 5 mL methanol was added 1.55 g (2.52 mmol) oxone followed by 500 μL water. The mixture was stirred at room temperature for 2 h, then partitioned between EtOAc and water, adding 1 N NaOH until the aqueous phase was neutral.

The organic layer was then dried over MgSO_4 and concentrated to obtain 387 mg product, 5-[N,N-(2-methylsulfonyl-ethyl)(4-fluorobenzyl)amino]-pyridin-2-yl methyl sulfone.

Example 12

5 2-[N,N-(2-methylsulfonyl-ethyl)(4-fluorobenzyl)amino]-pyridin-5-yl methyl sulfone (1-61)



Step 1

10 To a solution of 16 mL of 10% H_2O_2 in 24 mL concentrated H_2SO_4 at 0°C was added a solution of 2.34 g of 2-amino-5-bromopyridine dropwise with stirring. The ice bath was then removed, and allowed to warm to room temperature. After stirring at room temperature for 5 h, the reaction mixture was poured over ice and 1.62 g of the precipitated product, 5-bromo-2-nitro-pyridine, collected by vacuum filtration.

15 Step 2

To a solution of 1.0 g (4.93 mmol) 5-bromo-2-nitro-pyridine, dissolved in 10 mL DMF was added 379 mg (5.42 mmol) of sodium thiomethoxide followed by 569 mg (0.493 mmol) of tetrakis(triphenylphosphine)palladium(0). The mixture was heated to 80°C for 2 h, cooled to room temperature, and partitioned between ethyl acetate and water, dried over MgSO_4 and
20 concentrated. Purification by column chromatography, eluting with ethyl acetate/hexane, provided 318 mg of product, 5-methylthio-2-nitro-pyridine, pure by ^1H NMR.

Step 3

To 318 mg (1.86 mmol) of 5-methylthio-2-nitro-pyridine, dissolved in 10 mL acetone was added 7.44 mL (7.44 mmol) of a 1M solution of TiCl_3 dissolved in HCl. The mixture was
25 stirred at room temperature for 20 min, partitioned with ethyl acetate and 1N NaOH until

neutral, dried over MgSO_4 and concentrated to provide 225 mg of product, 2-amino-5-methylthio-pyridine, pure by ^1H NMR.

Step 4

To 112 μL (1.05 mmol) of 4-fluorobenzaldehyde and 147 mg (1.05 mmol) of 2-amino-5-methylthio-pyridine, dissolved in 6 mL dichloromethane was added 334 mg (1.57 mmol) sodium triacetoxyborohydride. The reaction mixture was stirred at room temperature for 4 h. The mixture was then partitioned between ethyl acetate and brine, dried over MgSO_4 and concentrated. Purification by column chromatography, eluting with ethyl acetate/hexane, provided 185 mg product, 2-[(4-fluorobenzyl)amino]-pyridin-5-yl methyl sulfide, pure by ^1H NMR.

Step 5


To 185 mg (0.744 mmol) of 2-[(4-fluorobenzyl)amino]-pyridin-5-yl methyl sulfide dissolved in 3 mL *N,N*-dimethylformamide was added 80 mg (0.744 mmol) methyl vinyl sulfone followed by 17 mg (0.744 mmol) sodium hydride. The mixture was stirred at room temperature for 0.25 h, partitioned between EtOAc and brine, dried over MgSO_4 and concentrated to obtain 262 mg of product, 2-[*N,N*-(2-methylsulfonyl-ethyl)(4-fluorobenzyl)amino]-pyridin-5-yl methyl sulfide.

Step 6

To 262 mg (0.74 mmol) of 2-[*N,N*-(2-methylsulfonyl-ethyl)(4-fluorobenzyl)amino]-pyridin-5-yl methyl sulfide dissolved in 3 mL methanol was added 939 mg (1.53 mmol) oxone followed by 500 μL water. The mixture was stirred at room temperature for 2 h, then partitioned between EtOAc and water, adding 1 N NaOH until the aqueous phase was neutral. The organic layer was then dried over MgSO_4 and concentrated to obtain 271 mg product of 2-[*N,N*-(2-methylsulfonyl-ethyl)(4-fluorobenzyl)amino]-pyridin-5-yl methyl sulfone.

Example 13

Synthesis of (4-ethoxy-benzyl)-(3-fluoro-4-methanesulfonyl-phenyl)-(2-methanesulfonyl-ethyl)-amine (1-72)



Step 1 (3-fluoro-4-thiomethyl-nitrobenzene):

A solution of 3,4-difluoronitrobenzene (5.0g) in 65mL dimethylformamide was treated all at once with sodium thiomethoxide (3.0g). After stirring overnight at ambient temperature under nitrogen the mixture was diluted with water and extracted with hexanes/ethyl acetate (1/1). The organic layers were combined and washed with water and brine. Filtration of the dried organic layer through a pad of silica gel was followed by solvent removal to afford 3.98g solid 3-fluoro-4-thiomethyl-nitrobenzene, which was carried on without characterization.

Step 2 (3-fluoro-4-thiomethyl-aniline):

A portion of the nitrobenzene from above (3.0g) was dissolved in 40mL of acetone and treated with titanium (III) chloride (50mL, 1.0M in HCl). After stirring 3h at ambient temperature under nitrogen the reaction was cautiously quenched with NaOH (1M aqueous solution), then with sodium bicarbonate (saturated aqueous solution). Extraction with three portions of ethyl acetate was performed, then the product was washed in the aqueous phase with three portions of 5% aqueous HCl. After making the aqueous washes basic with excess aqueous NaOH (1M), the product was washed into ethyl acetate. After drying, filtration and

solvent removal, 1.6g of 3-fluoro-4-thiomethyl-aniline was recovered as an oil. This material was used without further purification or characterization.

Step 3 ((4-ethoxy-benzyl)-(3-fluoro-4-methylthio-phenyl)-amine):

The aniline from above (0.51g, 3.24mmol) and 4-ethoxybenzaldehyde (0.50g, 3.33mmol) was dissolved in 1,2-dichloroethane (3 mL). Five drops of glacial acetic acid were added, followed by addition of sodium triacetoxyborohydride (1.2g, 5.7mmol). After stirring at ambient temperature over a weekend, the solution was directly poured onto a pad of silica gel and eluted with 20% ethyl acetate in hexanes. Solvent removal from the product containing fractions afforded the desired (4-ethoxy-benzyl)-(3-fluoro-4-methylthio-phenyl)-amine (1.0g, contaminated with 4-ethoxybenzaldehyde as indicated by ¹H NMR) as an oil, which was carried on to the final product as detailed below.

Step 4 ((4-ethoxy-benzyl)-(3-fluoro-4-methanesulfonyl-phenyl)-(2-methanesulfonyl-ethyl)-amine):

(4-ethoxy-benzyl)-(3-fluoro-4-methylthio-phenyl)-amine from above (0.40g) in 10 mL DMF was treated with methyl vinyl sulfone (0.35g), followed by sodium hydride (0.12g, 60% dispersion in mineral oil). After 3h stirring at ambient temperature under nitrogen, the reaction was quenched with aqueous sodium bicarbonate and extracted with ethyl acetate (two extractions). The combined organic layers were washed with brine, dried over magnesium sulfate, filtered, and solvent was removed. The residue was then diluted with 17mL of methanol and 3mL of water were added. Oxone (1.6g) was then added to the cooled solution (0°C) and the reaction was allowed to warm to room temperature. After stirring overnight, the reaction was diluted with water and extracted two times with ethyl acetate. The combined organic layers dried over magnesium sulfate, filtered, and solvent was removed. Flash chromatography starting with 33% ethyl acetate in hexanes, changing to 50% ethyl acetate in hexanes and finally 66% ethyl acetate in hexanes afforded, after solvent removal, 0.29g of the final product as a solid.


mp 56 – 59 C. Calc. for C₁₉H₂₄FO₅S₂ C 53.13 H 5.63 N 3.26. Found C 52.81 H 5.70 N 3.28.

Following the procedure above, but replacing 4-ethoxybenzaldehyde with 4-fluorobenzaldehyde in Step 3 gave (4-fluoro-benzyl)-(3-fluoro-4-methanesulfonyl-phenyl)-(2-methanesulfonyl-ethyl)-amine (1-73).

Following the procedure above, but replacing 3,4-difluoro-nitrobenzene with 2-bromo-5-nitroanisole in Step1 and 4-ethoxybenzaldehyde with 4-fluorobenzaldehyde in Step 3 gave (4-fluoro-benzyl)-(4-methanesulfonyl-3-methoxy-phenyl)-(2-methanesulfonyl-ethyl)-amine (1-74).

Example 14

4-[(2-methylsulfonyl-ethyl)-(2-methoxy-benzyl)-amino]-benzenesulfonamide (2-9)



5e

Step 1

A mixture of 4-fluorophenylsulfonamide (1.4g) and (2-thiomethyl)ethylamine (3 g) was heated at 120°C under a nitrogen atmosphere for 4 h; the mixture was then heated at 160°C for 2 h. The resulting dark mixture was cooled, passed through a pad of SiO₂ (hexane/EtOAc), to afford 0.34 g of 4-[(2-thiomethyl-ethyl)amino]-benzenesulfonamide as a white powder; ¹H NMR (DMSO) δ 2.1 (s, 3H), 2.65 (m, 2H), 3.3 (m, 2H), 6.5 (t, 1H, J = 5.8 Hz), 6.6 (m, 2H), 6.9 (s, 2H), 7.5 (m, 2H).

Step 2

A slurry of 4-[(2-thiomethyl-ethyl)amino]-benzenesulfonamide (0.33 g, 1.34 mmol) and 2-methoxybenzoyl chloride (0.22 mL, 0.25 g, 1.5 mmol) in (CH₂Cl)₂ was heated at reflux for 1 h. The mixture was evaporated *in vacuo*, and the residue was purified by MPLC (85:15 to 70:30 CH₂Cl₂/EtOAc) to afford 0.51 g (100%) of 4-[(2-methylthio-ethyl)-(2-methoxy-benzoyl)-amino]-benzenesulfonamide as a colorless glass; ¹H NMR δ 2.09 (s, 3H), 2.70 (m, 2H), 3.62 (s,

3H), 4.1 (m, 2H), 5.25 (s, 2H), 6.67 (m, 1H), 6.86 (t, 1H, J = 7.1), 7.2 (m, 4H), 7.72 (d, 2H, J = 7.1).

Step 4

5 A solution of 1 M BH₃•THF/THF (6.8 mL, 6.8 mmol) was added to a solution of 4-[(2-methylthio-ethyl)-(2-methoxy-benzoyl)-amino]-benzenesulfonamide (0.51 g, 1.34 mmol) in THF (5 mL). After 18 h, the excess BH₃ was quenched by addition of 0.1 M HCl; followed by partitioning the mixture between CH₂Cl₂ and NaHCO₃. The organic layer was dried (Na₂SO₄), filtered, evaporated *in vacuo*, and the residue was purified by MPLC (CH₂Cl₂ to 70:30
10 CH₂Cl₂/EtOAc) to afford 0.38 g (77%) of 4-[(2-methylthio-ethyl)-(2-methoxy-benzyl)-amino]-benzenesulfonamide as a clear glass.

Step 5

15 A solution of Oxone® (1.6 g, 2.6 mmol) in H₂O (5 mL) was added to a 0°C solution of 4-[(2-methylthio-ethyl)-(2-methoxy-benzyl)-amino]-benzenesulfonamide (0.38 g, 1.0 mmol) in MeOH (21 mL) resulting in an immediate precipitate. After 1 h, the mixture was partitioned between CH₂Cl₂ and H₂O. The aqueous layer was extracted with CH₂Cl₂ (2X). The combined organic layer was dried (Na₂SO₄), filtered, evaporated *in vacuo*, and the residue was triturated with hot CH₂Cl₂ to afford, after cooling, 0.38 g (92%) of 4-[(2-methylsulfonyl-ethyl)-(2-methoxy-benzyl)-amino]-benzenesulfonamide as a white solid; (m+H)⁺ 399.
20

Following the procedure of Example 14, but replacing 2-methoxybenzoyl chloride in Step 2 with 4-fluorobenzoyl chloride, 4-[(2-methylthio-ethyl)-(4-fluoro-benzoyl)-amino]-benzenesulfonamide was produced as a white solid; ¹H NMR δ 2.16 (s, 3H), 2.77 (m, 2H), 4.14 (m, 2H), 4.8 (br s, 2H), 6.90 (dd, 2H, J = 8.6, 8.6), 7.21 (d, 2H, J = 8.6), 7.32 (dd, 2H, J = 5.3, 8.9), 7.81 (d, 2H, J = 8.5) and 4-[(2-methylthio-ethyl)-(4-fluoro-benzyl)-amino]-benzenesulfonamide (**2-5**), was produced as a white solid; ¹H NMR δ 2.16 (s, 3H), 2.74 (m, 2H), 3.68 (m, 2H), 4.6 (br s, 2H), 6.1 (br s, 2H), 6.67 (d, 2H, J = 9.1), 7.01 (dd, 2H, J = 8.7, 8.7), 7.15 (dd, 2H, J = 5.3, 8.8), 7.70 (d, 2H, J = 9.1).

Following the procedure of Example 14, but replacing 2-methoxybenzoyl chloride in Step 2 with 4-ethoxybenzoyl chloride, 4-[(2-methylsulfonyl-ethyl)-(4-ethoxy-benzyl)-amino]-benzenesulfonamide, (**2-6**), was produced as a white solid; (m+H)⁺ 413.


Following the procedure of Example 14, but replacing 2-methoxybenzoyl chloride in Step 2 with 2-fluorobenzoyl chloride, 4-[(2-methylsulfonyl-ethyl)-(2-fluoro-benzyl)-amino]-benzenesulfonamide, (**2-7**), was produced as a white solid; (m+H)⁺ 387.

Following the procedure of Example 14, but replacing 2-methoxybenzoyl chloride in Step 2 with 2,6-difluorobenzoyl chloride, 4-[(2-methylsulfonyl-ethyl)-(2,6-difluoro-benzyl)-amino]-benzenesulfonamide, (**2-8**), was produced as a white solid; (m+H)⁺ 405.

Following the procedure of Example 14, but replacing 2-methoxybenzoyl chloride in Step 2 with 2-chlorobenzoyl chloride, 4-[(2-methylsulfonyl-ethyl)-(2-chloro-benzyl)-amino]-benzenesulfonamide, (**2-10**), was produced as a white solid; (m+H)⁺ 403.

Example 15

4-[(2-methylsulfonyl-ethyl)-(4-fluoro-benzyl)amino]-N-ethyl-benzenesulfonamide (**2-12**)



Step 1

A solution of 4-[(2-methylthio-ethyl)-(4-fluoro-benzoyl)-amino]-benzenesulfonamide (0.32 g, 0.88 mmol), acetyl chloride (0.069 mL, 76 mg, 0.97 mmol), and Et₃N (0.13 mL, 97 mg, 0.96 mmol) in CH₂Cl₂ (9 mL) was heated at reflux for 2 h. After cooling, the mixture was evaporated *in vacuo*, and the residue was purified by MPLC (CH₂Cl₂ to 60:40 CH₂Cl₂/EtOAc) to afford 0.29 g (81%) of 4-[(2-methylthio-ethyl)-(4-fluoro-benzoyl)-amino]-N-acetyl-benzenesulfonamide as a colorless glass; ¹H NMR δ 2.03 (s, 3H), 2.15 (s, 3H), 2.77 (m, 2H),

4.15 (m, 2H), 6.90 (dd, 2H, J = 7.5, 7.5), 7.24 (d, 2H, J = 9.0), 7.32 (dd, 2H, J = 3.0, 9.0), 7.92 (d, 2H, J = 9.0), 9.1 (br s, 1H).

Step 2

5 A solution of 1 M $\text{BH}_3 \cdot \text{THF}$ (7.0 mL, 7.0 mmol) was added to a stirring solution of 4-[(2-methylthio-ethyl)-(4-fluoro-benzoyl)-amino]-N-acetyl-benzenesulfonamide (0.29 g, 0.69 mmol) in THF (8 mL). After 18 h, the excess BH_3 was quenched by addition of 0.1 M HCl; followed by partitioning the mixture between CH_2Cl_2 and NaHCO_3 . The organic layer was dried (Na_2SO_4), filtered, evaporated *in vacuo*, and the residue was purified by MPLC
10 (CH_2Cl_2 to 90:10 $\text{CH}_2\text{Cl}_2/\text{EtOAc}$) to afford 0.16 g (59%) of 4-[(2-methylthio-ethyl)-(4-fluoro-benzyl)-amino]-N-ethyl-benzenesulfonamide as a colorless glass; ^1H NMR δ 1.09 (t, 3H, J = 7.5), 2.16 (s, 3H), 2.74 (m, 2H), 2.96 (m, 2H), 3.67 (m, 2H), 4.37 (t, 1H, J = 6.0), 4.62 (s, 2H), 6.69 (d, 2H, J = 9.0), 7.04 (dd, 2H, J = 4.5, 7.5), 7.15 (dd, 2H, J = 6.0, 9.0), 7.66 (d, 2H, J = 7.5).


Step 3

15 A solution of Oxone® (0.54 g, 0.88 mmol) in H_2O (2 mL) was added to a 0°C solution of 4-[(2-methylthio-ethyl)-(4-fluoro-benzyl)-amino]-N-ethyl-benzenesulfonamide (0.13 g, 0.34 mmol) in MeOH (8 mL) resulting in an immediate precipitate. After 1 h, the mixture was
20 partitioned between CH_2Cl_2 and H_2O . The aqueous layer was extracted with CH_2Cl_2 (2X). The combined organic layer was dried (Na_2SO_4), filtered, and evaporated *in vacuo* to afford 0.14 g (100%) of 4-[(2-methylsulfonyl-ethyl)-(4-fluoro-benzyl)-amino]-N-ethyl-benzenesulfonamide as a pale glass; $(\text{m}+\text{H})^+$ 415.

Example 16

4-[(2-methylsulfonyl-ethyl)-(2-fluoro-benzyl)amino]-(N-2-fluorobenzyl)-benzenesulfonamide
(2-11)

5




Step 1

A solution of 4-[(2-thiomethyl-ethyl)amino]-benzenesulfonamide (0.30 g, 1.2 mmol), 2-fluorobenzoyl chloride (0.29 mL, 0.39 g, 2.4 mmol), and Et₃N (0.34 mL, 0.25 g, 2.4 mmol) in CH₂Cl₂ (12 mL) was heated at reflux for 2 h. After cooling, the mixture was evaporated *in vacuo*, and the residue was purified by MPLC (CH₂Cl₂ to 75:25 CH₂Cl₂/EtOAc) to afford 0.20 g (33%) of 4-[(2-methylthio-ethyl)-(2-fluoro-benzoyl)-amino]-(N-2-fluorobenzoyl)-benzenesulfonamide 3g as a colorless glass; ¹H NMR δ 2.13 (s, 3H), 2.74 (m, 2H), 4.13 (m, 2H), 6.80 (t, 1H, J = 9.0), 7.06 (t, 1H, J = 7.5), 7.17 (dd, 1H, J = 7.5, 12.0), 7.2-7.3 (m, 4H), 7.36 (t, 1H, J = 7.5), 7.59 (m, 1H), 7.97 (t, 1H, J = 7.5), 8.02 (d, 2H, J = 9.0), 9.02 (d, 1H, J = 15.0).

Following the procedure of Example 15, Step 2, but replacing 4-[(2-methylthio-ethyl)-(4-fluoro-benzoyl)-amino]-N-acetyl-benzenesulfonamide with 4-[(2-methylthio-ethyl)-(2-fluoro-benzoyl)-amino]-(N-2-fluorobenzoyl)-benzenesulfonamide, was produced 4-[(2-methylsulfonyl-ethyl)-(2-fluoro-benzyl)-amino]-N-(2-fluoro-benzyl)-benzenesulfonamide as a pale glass; (m+H)⁺ 495.

Example 17


2-Fluoro-5-[[[(4-methanesulfonyl-phenyl)-(3-methanesulfonyl-propyl)-amino]-methyl]-phenol (1-79)



To 694 mg (1.67 mmol) of (4-fluoro-3-methoxy-benzyl)-(4-methanesulfonyl-phenyl)-(3-methanesulfonyl-propyl)-amine, prepared following the method of Example 5 but replacing 3-pyridinecarboxaldehyde with 4-fluoro-3-methoxybenzaldehyde, dissolved in 3 mL of 2,4,6-collidine was added 402 mg (3.01 mmol) lithium iodide. The mixture was heated to 150 degrees for 3 hrs, cooled to room temperature, and partitioned between ethyl acetate and 1 N HCl. Upon drying over magnesium sulfate and concentration, column chromatography, eluting with acetone/dichloromethane, provided 169 mg of 2-fluoro-5-[[[(4-methanesulfonyl-phenyl)-(3-methanesulfonyl-propyl)-amino]-methyl]-phenol; (m+H)⁺= 401.

Example 18

Synthesis of (4-ethoxy-benzyl)-(4-methanesulfonyl-phenyl)-thiophen-3-ylmethyl-amine (1-76)



Step 1

To 5.0 mL (40.19 mmol) of 4-(methylthio)aniline dissolved in 25 mL dichloromethane was added 5.59 mL (40.19 mmol) 4-ethoxybenzaldehyde followed by 12.78 g (60.28 mmol) sodium triacetoxyborohydride. The mixture was stirred overnight at room temperature, partitioned between EtOAc and brine, dried over MgSO₄ and concentrated. Crystallization from dichloromethane/hexane, provided 7.87 g of (4-Ethoxy-benzyl)-(4-methylsulfanyl-phenyl)-amine, pure by ¹H NMR.

Step 2

To 200 mg (0.731 mmol) of (4-Ethoxy-benzyl)-(4-methylsulfanyl-phenyl)-amine dissolved in 5 mL dichloromethane was added 68 µL (0.731 mmol) 3-thiophenecarboxaldehyde followed by 232 mg (1.09 mmol) sodium triacetoxyborohydride. The mixture was stirred overnight at room temperature, partitioned between EtOAc and brine, dried over MgSO₄ and concentrated. Column chromatography, eluting with ethyl acetate/hexane, provided 241 mg product (4-Ethoxy-benzyl)-(4-methylsulfanyl-phenyl)-thiophen-3-ylmethyl-amine, pure by ¹H NMR.

Step 3

To 241 mg (0.652 mmol) of (4-Ethoxy-benzyl)-(4-methylsulfanyl-phenyl)-thiophen-3-ylmethyl-amine dissolved in 6 mL methanol was added 800 mg (1.3 mmol) OXONETM followed by 600 µL water. The mixture was stirred at room temperature for 2 h, then partitioned between EtOAc and water, adding 1 N NaOH until the aqueous phase was neutral. The organic layer was then dried over MgSO₄ and concentrated. (4-Ethoxy-benzyl)-(4-methanesulfonyl-phenyl)-thiophen-3-ylmethyl-amine was obtained in 92% yield (240 mg) and appeared pure by ¹H NMR.

Following the procedure of Example 18, but replacing 2-thiophenecarboxaldehyde in step 2 with 4-imidazolecarboxaldehyde gave (4-Ethoxy-benzyl)-(1*H*-imidazol-4-ylmethyl)-(4-methanesulfonyl-phenyl)-amine. (m+H)⁺= 385. (1-77)

Example 19

The following are representative pharmaceutical formulations containing a compound of formula (I).

Tablet formulation

5 The following ingredients are mixed intimately and pressed into single scored tablets.

| | Ingredient | Quantity per tablet, mg |
|----|----------------------------|----------------------------|
| 10 | compound of this invention | 400 |
| | cornstarch | 50 |
| | croscarmellose sodium | 25 |
| | lactose | 120 |
| | magnesium stearate | 5 |

Capsule formulation

15 The following ingredients are mixed intimately and loaded into a hard-shell gelatin capsule.

| | Ingredient | Quantity per capsule, mg |
|----|----------------------------|-----------------------------|
| 20 | compound of this invention | 200 |
| | lactose, spray-dried | 148 |
| | magnesium stearate | 2 |

Suspension formulation

25 The following ingredients are mixed to form a suspension for oral administration.

| | Ingredient | Amount |
|----|----------------------------|----------------|
| | compound of this invention | 1.0 g |
| | fumaric acid | 0.5 g |
| | sodium chloride | 2.0 g |
| 30 | methyl paraben | 0.15 g |
| | propyl paraben | 0.05 g |
| | granulated sugar | 25.5 g |
| | sorbitol (70% solution) | 12.85 g |
| | Veegum K (Vanderbilt Co.) | 1.0 g |
| 35 | flavoring | 0.035 ml |
| | colorings | 0.5 mg |
| | distilled water | q.s. to 100 ml |

Injectable formulation

The following ingredients are mixed to form an injectable formulation.

| 5 | <table><tr><th>Ingredient</th><th>Amount</th></tr><tr><td>compound of this invention</td><td>0.4 mg</td></tr><tr><td>sodium acetate buffer solution, 0.4 M</td><td>2.0 ml</td></tr><tr><td>HCl (1N) or NaOH (1N)</td><td>q.s. to suitable pH</td></tr><tr><td>water (distilled, sterile)</td><td>q.s. to 20 ml</td></tr></table> | Ingredient | Amount | compound of this invention | 0.4 mg | sodium acetate buffer solution, 0.4 M | 2.0 ml | HCl (1N) or NaOH (1N) | q.s. to suitable pH | water (distilled, sterile) | q.s. to 20 ml |
|---------------------------------------|--|------------|--------|----------------------------|--------|---------------------------------------|--------|-----------------------|---------------------|----------------------------|---------------|
| Ingredient | Amount | | | | | | | | | | |
| compound of this invention | 0.4 mg | | | | | | | | | | |
| sodium acetate buffer solution, 0.4 M | 2.0 ml | | | | | | | | | | |
| HCl (1N) or NaOH (1N) | q.s. to suitable pH | | | | | | | | | | |
| water (distilled, sterile) | q.s. to 20 ml | | | | | | | | | | |

10

Example 20

Inhibition of COX I and COX II *in vitro*

The COX I and COX II inhibitory activity of compounds of this invention *in vitro* was
15 determined using partially purified COX I and COX II enzymes, prepared as described in J.
Barnett *et. al.*, *Biochim. Biophys. Acta*, **1209**, 130-139 (1994).

COX I and COX II samples were diluted with Tris-HCl buffer (50mM Tris-HCl, pH 7.9)
containing 2 mM EDTA and 10% glycerol and reconstituted by incubating first with 2 mM
phenol for 5 minutes and then with 1 micromolar hematin for an additional 5 minutes. 125 µl of
20 the reconstituted COX I or COX II enzyme were preincubated for 10 minutes at room temperature
in a shaking water bath with the compounds of the invention dissolved in 2-15 µl of DMSO or
the carrier vehicles (control samples). The enzyme reaction was initiated by adding 25 µl of 1-[14
C]arachidonic acid (80,000-100,000 cpm/tube; 20 micromolar final concentration) and the
reaction was allowed to continue for an additional 45 seconds. The reaction was terminated by
25 adding 100 µl of 2N HCl and 750 µl water. An aliquot (950 µl) of the reaction mixture was
loaded onto a 1 ml C₁₈ Sep-Pak column (J.T. Baker, Phillipsburg, NJ) which had been previously
washed with 2-3 ml methanol and equilibrated with 5-6 ml distilled water. Oxygenated products
were quantitatively eluted with 3 ml of acetonitrile/water/acetic acid (50:50:0.1, v/v) and the
radioactivity in the eluate determined in a scintillation counter.
30 Compounds of this invention were active in this assay.

The COX inhibitory activities (expressed as IC₅₀ , the concentration causing 50% inhibition of the COX enzyme being assayed) of some compounds of the invention were:

| CPD # | COX I IC ₅₀ , μM | COX II IC ₅₀ , μM | CPD # | COX I IC ₅₀ , μM | COX II IC ₅₀ , μM |
|-------|-----------------------------|------------------------------|-------|-----------------------------|------------------------------|
| 1-24 | >40 | <0.20 | 1-48 | >15 | <0.20 |
| 1-26 | >40 | <0.20 | 1-49 | >40 | <0.20 |
| 1-27 | | <0.20 | 1-50 | >40 | <0.20 |
| 1-37 | >40 | <0.20 | 1-51 | >40 | <0.20 |
| 1-47 | >20 | <0.20 | 1-53 | >40 | <0.20 |

5

Example 21

Anti-inflammatory activity

The anti-inflammatory activity of compounds of this invention may be determined by measuring the inhibition of carrageenan-induced paw edema in the rat, using a modification of the method described in Winter C. A. *et al.*, "Carrageenan-Induced Edema in Hind Paw of the Rat as an Assay for Anti-inflammatory Drugs" *Proc. Soc. Exp. Biol. Med.* **111**, 544-547, (1962). This assay has been used as a primary *in vivo* screen for anti-inflammatory activity of most NSAIDs, and is considered predictive of human efficacy. Briefly, test materials are administered orally to female rats in a volume of 1 ml prepared as solutions or suspensions in an aqueous vehicle containing 0.9% sodium chloride, 0.5% sodium carboxymethyl-cellulose, 0.4% polysorbate 80, 0.9% benzyl alcohol and 97.3% distilled water. Control rats receive vehicle alone. After 1 h 0.05 ml of a 0.5% solution of Carrageenan (Type IV Lambda, Sigma Chemical Co.) in 0.9% saline is injected into the subplantar region of the right hind paw. Three hours later the rats are euthanized in a carbon dioxide atmosphere; hind paws are removed by severing at the tarso-crural joint; and the left and right paws are weighed. The increase in weight of the right paw over the left paw is obtained for each animal and the mean increases are calculated for each group. The anti-inflammatory activity of the test materials is expressed as the percent inhibition of the increase in hind paw weight of the test group relative to the vehicle dosed control group.

Compounds of the invention are active in this assay.

Example 22

Inhibition of eicosanoid synthesis *in vivo*

5 The activity of compounds of this invention in inhibiting *in vivo* eicosanoid (prostaglandin E₂) synthesis in inflamed tissues may be determined by the carrageenan-induced inflammation (air-pouch model) in rats, using a modification of the method described in Futaki, M., *et al.*, "Selective Inhibition of NS-398 on prostanoid production in inflamed tissue in rat Carrageenan Air-pouch Inflammation" *J. Pharm. Pharmacol.* **45**, 753-755, (1993) and Masferrer, J. L., *et al.*;

10 "Selective Inhibition of inducible cyclooxygenase 2 *in vivo* is Antiflammatory and Nonulcerogenic" *Proc. Natl. Acad. Sci. USA.* **91**, 3228-3232, (1994). In this assay, an air-pouch is created in the rat and the PGE₂ levels in the air-pouch exudate are measured by enzyme immunoassay. Briefly, male rats are anesthetized using a 60:40 CO₂:O₂ mixture and subsequently injected subcutaneously with 20 ml of sterilized air, under aseptic conditions, in the proximal area

15 of the dorsum. This injection of sterile air causes the creation of a subcutaneous "air pouch". The next day, a further 10 ml of sterile air is injected into the previously formed pouch using the same technique. The test materials are administered orally in a volume of 1ml/100g body weight as solutions or suspensions in an aqueous vehicle containing 0.9% sodium chloride, 0.5% sodium carboxymethyl-cellulose, 0.4% polysorbate 80, 0.9% benzyl alcohol and 97.3% water. Control

20 rats receive vehicle alone. After 30 minutes, 5 ml of a 0.5% solution of carrageenan (Sigma, Lambda Type IV) is injected into the air pouch. The rats are euthanized 3 or 6 h after the compound administration. 10 ml of a solution containing 10 µg/l of indomethacin and 5.4 mM EDTA in 0.9% sterile saline is injected into the air pouch; the air pouch is cut open; and the exudate is harvested. The total exudate volume is recorded, and the samples are analyzed for

25 PGE₂ and 6-keto PGF₁ by ELISA (Titerzyme[®], PerSeptive Diagnostics, Boston, MA) and TxB₂ by radioimmuno assay (New England Nuclear Research, Boston MA, Catalog No. NEK-037), according to the manufacturer's directions.

 The mean concentrations of PGE₂ are calculated for each group. The anti-inflammatory activity of test materials is expressed as the percent inhibition of PGE₂ formation in the test group

30 relative to the control group.

Example 23

Analgesic Activity

The analgesic activity of the compounds of this invention may be determined by using a modification of the method described in Randall, L. O., and Selitto, J. J., “ A Method for

5 Measurement of Analgesic Activity on Inflamed Tissue”, *Arch. Int. Pharmacodyn.*, CXI, 4, 409, (1957) and Gans, *et. al.*, “ Anti-Inflammatory and Safety Profile of DuP 697, a Novel Orally Effective Prostaglandin Synthesis Inhibitor”, *J. Pharmacol. Exp. Ther.*, **254**, No. 1, 180, (1990). In this assay, the male Sprague Dawley rats are injected with 0.1 ml of 20% brewer’s yeast in deionized water (Sigma, St. Louis) in the subplantar region of the left hind foot. After
10 2 h, the test materials are administered orally in a volume of 1 ml/100g body weight as solutions or suspensions in an aqueous vehicle containing 0.9% sodium chloride, 0.5% sodium carboxymethyl-cellulose, 0.4% polysorbate 80, 0.9% benzyl alcohol and 97.3% water. Control rats receive vehicle alone. After 1 h, the hindpaw is placed on the platform of a Basile Analgesy-Meter (Ugo Biological Research Apparatus, Italy, Model # 7200) and mechanical
15 force is applied to the dorsum of the rat’s hindpaw. The analgesic activity of compounds of this invention may also be determined by using an adjuvant-induced arthritis pain model in the rat, where pain is assessed by the animal's vocal response to the squeezing or flexing of an inflamed ankle joint, as described in Winter C.A. and Nuss, G.W., “Treatment of Adjuvant Arthritis in rats with Antiinflammatory Drugs”, *Arthritis Rheum.*, **9**, 394-403, (1966) and Winter, C.A.,
20 Kling P.J., Tocco, D.J., and Tanabe, K., “Analgesic activity of Diflunisal [MK-647; 5-(2,4-Difluorophenyl)salicylic acid] in Rats with Hyperalgesia Induced by Freund's Adjuvant”, *J. Pharmacol. Exp. Ther.*, **211**, 678-685, (1979).

The foregoing invention has been described in some detail by way of illustration and
25 example, for purposes of clarity and understanding. It will be obvious to one of skill in the art that changes and modifications may be practiced within the scope of the appended claims. Therefore, it is to be understood that the above description is intended to be illustrative and not restrictive. The scope of the invention should, therefore, be determined not with reference to the above description, but should instead be determined with reference to the following
30 appended claims, along with the full scope of equivalents to which such claims are entitled.

